

527 Rec'd PCT/PTO 29 NOV 2000

CERTIFICATION UNDER 37 CFR 1.10

09/701500

EI685061565US

Express Mail Mailing Number

29 NOV 2000

Date of Deposit

I hereby certify that this paper or fee, and any documents referred to as enclosed herein, are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Box PCT, Washington, D.C. 20231.

Cederic Rodgers

Name of Person Mailing Application

Signature of Person Mailing Application

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TSRI 651.1

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)
Not known yet

INTERNATIONAL APPLICATION NO.
PCT/US99/11780

INTERNATIONAL FILING DATE
28 MAY 1999

PRIORITY DATE CLAIMED
29 MAY 1998

TITLE OF INVENTION

METHODS AND COMPOSITIONS USEFUL FOR MODULATION OF ANGIOGENESIS USING TYROSINE KINASE SRC

APPLICANT(S) FOR DO/EO/US

David A. Cheresch, Brian Eliceiri and Pamela L. Schwartzberg

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ The US has been elected by the expiration of 19 months from the priority date (PCT Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☐ An English language translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) 35 U.S.C. 371(c)(4)) and **Power of Attorney** (Executed).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 16 below concern other document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A FIRST preliminary amendment.
- ☐ A SECOND or SUBSEQUENT preliminary amendment.

545 Rec'd PCT/PTO 29 NOV 2009

| | | |
|--|---|---------------------------------------|
| US APPLICATION NO. (if known, see 37 CFR 1.5) <div style="font-size: 1.5em; font-weight: bold; margin-top: 10px;">09/701500</div> | INTERNATIONAL APPLICATION NO. PCT/US99/11780 | ATTORNEYS DOCKET NUMBER TSRI 651.1 |
|--|---|---------------------------------------|

| | |
|---|--|
| 14. <input type="checkbox"/> A substitute specification and claims amended to conform the translated application to US practice 15. <input type="checkbox"/> A change of power of attorney and/or address letter. 16. <input checked="" type="checkbox"/> Other items or information: (a) copy of form PCT/IB/308 (Notification of Communication of International Application to the Designated Offices); (b) copy of the First Page of International Publication; (c) copy of International Search Report; (d) copies of three (3) forms PCT/IB/306 (Notification of the Recording of a Change); (e) copies of Written Opinion and International Preliminary Examination Report | |
|---|--|

| 17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492(A)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1,000.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 <div style="text-align: right;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div> | CALCULATIONS PTO USE ONLY | | | | | | | | | | | | | | | | |
|--|---|--------------|--------------|------|--------------|----|----|---------|--------------------|---|---|---------|---|--|--|----------|---|
| Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)). | <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;">\$690.00</div> <div style="border: 1px solid black; padding: 5px;">\$0.00</div> | | | | | | | | | | | | | | | | |
| <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 20%;">Claims</th> <th style="width: 20%;">Number Filed</th> <th style="width: 20%;">Number Extra</th> <th style="width: 20%;">Rate</th> </tr> <tr> <td>Total Claims</td> <td style="text-align: center;">38</td> <td style="text-align: center;">18</td> <td style="text-align: center;">\$18.00</td> </tr> <tr> <td>Independent Claims</td> <td style="text-align: center;">8</td> <td style="text-align: center;">5</td> <td style="text-align: center;">\$80.00</td> </tr> <tr> <td colspan="3">MULTIPLE DEPENDENT CLAIM(S) (if applicable)</td> <td style="text-align: center;">\$270.00</td> </tr> </table> | Claims | Number Filed | Number Extra | Rate | Total Claims | 38 | 18 | \$18.00 | Independent Claims | 8 | 5 | \$80.00 | MULTIPLE DEPENDENT CLAIM(S) (if applicable) | | | \$270.00 | <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;">\$324.00</div> <div style="border: 1px solid black; padding: 5px;">\$400.00</div> |
| Claims | Number Filed | Number Extra | Rate | | | | | | | | | | | | | | |
| Total Claims | 38 | 18 | \$18.00 | | | | | | | | | | | | | | |
| Independent Claims | 8 | 5 | \$80.00 | | | | | | | | | | | | | | |
| MULTIPLE DEPENDENT CLAIM(S) (if applicable) | | | \$270.00 | | | | | | | | | | | | | | |
| TOTAL OF ABOVE CALCULATIONS = | <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;">\$1,414.00</div> <div style="border: 1px solid black; padding: 5px;">\$0.00</div> | | | | | | | | | | | | | | | | |
| <input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2. | <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;">\$1,414.00</div> <div style="border: 1px solid black; padding: 5px;">\$0.00</div> | | | | | | | | | | | | | | | | |
| SUBTOTAL = | <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;">\$1,414.00</div> <div style="border: 1px solid black; padding: 5px;">\$0.00</div> | | | | | | | | | | | | | | | | |
| Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)). | <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;">\$0.00</div> <div style="border: 1px solid black; padding: 5px;">\$0.00</div> | | | | | | | | | | | | | | | | |
| TOTAL NATIONAL FEE = | <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;">\$1,414.00</div> <div style="border: 1px solid black; padding: 5px;">\$0.00</div> | | | | | | | | | | | | | | | | |
| Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) \$40 per property | <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;">\$1,414.00</div> <div style="border: 1px solid black; padding: 5px;">\$0.00</div> | | | | | | | | | | | | | | | | |
| TOTAL FEES ENCLOSED = | <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;">\$1,414.00</div> <div style="border: 1px solid black; padding: 5px;">\$0.00</div> | | | | | | | | | | | | | | | | |
| | Amount to be refunded \$ | | | | | | | | | | | | | | | | |
| | charged \$ | | | | | | | | | | | | | | | | |

09701500 112900
525 Rec'd PCT/PTO 29 NOV 2000

| | | |
|---|---|---------------------------------------|
| US APPLICATION NO. (if known, see 37 CFR 1.101) 09/701500 | INTERNATIONAL APPLICATION NO. PCT/US99/11780 | ATTORNEYS DOCKET NUMBER TSRI 651.1 |
|---|---|---------------------------------------|


- a. ☒ A check in the amount of \$1,414.00 to cover the above fees is enclosed.
- b. ☐ Please charge my Deposit Account No. _____ in the amount of _____ to cover the above fees. A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional filing fees which may be required, or credit any overpayment to Deposit Account No. 15-0508. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

OLSON & HIERL, LTD.
20 North Wacker Drive
36th Floor
Chicago, Illinois 60606

Telephone: (312) 580-1180
Customer No.: 002387



SIGNATURE

Talivaldis Cepuritis

NAME

20,818

REGISTRATION NO.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US)

Applicant(s): David A. Cheresh et al.)
Serial No.: 09/701,500)
Filed: November 29, 2000)
For: METHODS AND COMPOSITIONS)
USEFUL FOR MODULATION OF)
ANGIOGENESIS USING)
TYROSINE KINZE SRC)
Atty. Docket No.: TSRI 651.1

COMMUNICATION

Box PCT
Commissioner for Patents
Washington, D.C. 20231

Sir:

Pursuant to the phone request of Mr. John Anderson of the U.S. Patent Office on November 14, 2002, enclosed is a duplicate diskette of the "Sequence Listing" to replace the diskette misplaced.

This submission does not constitute new matter and is supported in the application as filed.

Respectfully submitted,

By Deborah L. Boklund-Moran
Deborah L. Boklund-Moran
Foreign Patent Paralegal for
Talivaldis Cepuritis (Reg. No. 20,818)

OLSON & HIERL, LTD.
20 North Wacker Drive
36th Floor
Chicago, Illinois 60606
(312) 580-1180

Serial No. 09/701,500 - - - - - 2

CERTIFICATION UNDER 37 CFR 1.10

"Express Mail" mailing label number: EL 667310310 US

I hereby certify that this correspondence, together with any other documents and/or fees referred to as enclosed herein, is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service on November 15, 2002 and is addressed to: Box PCT, Commissioner for Patents, Washington, D.C. 20231.

Cedric Rodgers

(Typed or printed name of person mailing paper or fee)

Cedric Rodgers

(Signature of person mailing paper or fee)

09701500.112500

#3

JC17 Rec'd PCT/PTO 24 MAY 2001

09/701500

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US)

Applicant(s): David A. Cheresh et al.)
Serial No.: 09/701,500)
Filed: November 29, 2000)
For: METHODS AND COMPOSITIONS)
USEFUL FOR MODULATION OF) Atty. Docket No.: TSRI 651.1
ANGIOGENESIS USING)
TYROSINE KINZE SRC)

COMMUNICATION


Box PCT
Commissioner for Patents
Washington, D.C. 20231

Sir:

In response to the Notification of Missing Requirements Under 35 U.S.C. 371 In the United States Designated/Elected Office (DO/EO/US) dated April 30, 2001, enclosed is a substitute paper copy and computer-readable form of the "Sequence Listing." A copy of the Patent Office Notification is returned herewith.

This submission does not constitute new matter and is supported in the application as filed. To the best of my information and belief, the sequence listing information recorded in computer-readable form is identical to the paper copy of the sequence listing.

Respectfully submitted,

By 
Talivaldis Cepuritis (Reg. No. 20,818)

OLSON & HIERL, LTD.
20 North Wacker Drive
36th Floor
Chicago, Illinois 60606
(312) 580-1180

Serial No. 09/701,500 - - - - - 2

CERTIFICATION UNDER 37 CFR 1.10

"Express Mail" mailing label number: EL 841695176 US

I hereby certify that this correspondence, together with any other documents and/or fees referred to as enclosed herein, is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service on May 24, 2001 and is addressed to: Box PCT, Commissioner for Patents, Washington, D.C. 20231.

Cedric Rodgers

(Typed or printed name of person mailing paper or fee)

Cedric Rodgers

(Signature of person mailing paper or fee)

METHODS AND COMPOSITIONS USEFUL FOR MODULATION
OF ANGIOGENESIS USING TYROSINE KINASE SRC

Reference to Related Application

5 This application claims the benefit of U.S. Provisional Application Serial
No. 60/087,220 filed May 29, 1998, which is incorporated by reference, as are all
references cited herein.

Technical Field

10 The present invention relates generally to the field of medicine, and relates
specifically to methods and compositions for modulating angiogenesis of tissues
using the protein tyrosine kinase Src, variants of Src, and nucleic acids encoding
them.

Background

15 Angiogenesis is a process of tissue vascularization that involves the growth of
new developing blood vessels into a tissue, and is also referred to as neo-
vascularization. The process is mediated by the infiltration of endothelial cells and
smooth muscle cells. The process is believed to proceed in any one of three ways:
the vessels can sprout from pre-existing vessels, de-novo development of vessels
can arise from precursor cells (vasculogenesis), or existing small vessels can
20 enlarge in diameter. Blood et al., Bioch. Biophys. Acta, 1032:89-118 (1990).

Angiogenesis is an important process in neonatal growth, but is also important
in wound healing and in the pathogenesis of a large variety of clinical diseases
including tissue inflammation, arthritis, tumor growth, diabetic retinopathy,
macular degeneration by neovascularization of the retina and like conditions.
25 These clinical manifestations associated with angiogenesis are referred to as
angiogenic diseases. Folkman et al., Science, 235:442-447 (1987). Angiogenesis
is generally absent in adult or mature tissues, although it does occur in wound
healing and in the corpus luteum growth cycle. See, for example, Moses et al.,
Science, 248:1408-1410 (1990).

30 It has been proposed that inhibition of angiogenesis would be a useful therapy
for restricting tumor growth. Inhibition of angiogenesis has been proposed by (1)
inhibition of release of "angiogenic molecules" such as bFGF (basic fibroblast
growth factor), (2) neutralization of angiogenic molecules, such as by use of anti-

β bFGF antibodies, (3) use of inhibitors of vitronectin receptor $\alpha_v\beta_3$, and (4) inhibition of endothelial cell response to angiogenic stimuli. This latter strategy has received attention, and Folkman et al., Cancer Biology, 3:89-96 (1992), have described several endothelial cell response inhibitors, including collagenase inhibitor, basement membrane turnover inhibitors, angiostatic steroids, fungal-derived angiogenesis inhibitors, platelet factor 4, thrombospondin, arthritis drugs such as D-penicillamine and gold thiomalate, vitamin D₃ analogs, alpha-interferon, and the like that might be used to inhibit angiogenesis. For additional proposed inhibitors of angiogenesis, see Blood et al., Bioch. Biophys. Acta., 1032:89-118 (1990), Moses et al., Science, 248:1408-1410 (1990), Ingber et al., Lab. Invest., 59:44-51 (1988), and United States Patent Nos. 5,092,885, 5,112,946, 5,192,744, 5,202,352, 5,753,230 and 5,766,591. None of the inhibitors of angiogenesis described in the foregoing references involve the Src proteins.

For angiogenesis to occur, endothelial cells must first degrade and cross the blood vessel basement membrane in a similar manner used by tumor cells during invasion and metastasis formation.

It has been previously reported that angiogenesis depends on the interaction between vascular integrins and extracellular matrix proteins. Brooks et al., Science, 264:569-571 (1994). Furthermore, it was reported that programmed cell death (apoptosis) of angiogenic vascular cells is initiated by the interaction, which would be inhibited by certain antagonists of the vascular integrin $\alpha_v\beta_3$. Brooks et al., Cell, 79:1157-1164 (1994). More recently, it has been reported that the binding of matrix metalloproteinase-2 (MMP-2) to vitronectin receptor ($\alpha_v\beta_3$) can be inhibited using $\alpha_v\beta_3$ antagonists, and thereby inhibit the enzymatic function of the proteinase. Brooks et al., Cell, 85:683-693 (1996).

Summary of the Invention

The present invention is directed to modulation of angiogenesis in tissues by tyrosine kinase Src, also referred to generically herein as Src.

Compositions and methods for modulating angiogenesis in a tissue associated with a disease condition are contemplated. A composition comprising an angiogenesis-modulating amount of a Src protein is administered to tissue to be treated for a disease condition that responds to modulation of angiogenesis. The composition providing the Src protein can contain purified protein, biologically

active protein fragments, recombinantly produced Src protein or protein fragments or fusion proteins thereof, or gene/nucleic acid expression vectors for expressing a Src protein.

Where the Src protein is inactivated or inhibited, the modulation is an inhibition of angiogenesis. Where the Src protein is active or activated, the modulation is a potentiation of angiogenesis.

The tissue to be treated can be any tissue in which modulation of angiogenesis is desirable. For angiogenesis inhibition, it is useful to treat diseased tissue where deleterious neovascularization is occurring. Exemplary tissues include inflamed tissue, solid tumors, metastases, tissues undergoing restenosis, and the like tissues.

For potentiation, it is useful to treat patients with ischemic limbs in which there is poor circulation in the limbs from diabetic or other conditions. Patients with chronic wounds that do not heal and therefore could benefit from the increase in vascular cell proliferation and neovascularization can be treated as well.

Particularly preferred is the use of Src protein containing a modified amino acid sequence as described herein. Several particularly useful modified Src proteins and the expression thereof are described herein.

The present invention also encompasses a pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a viral or non-viral gene transfer vector containing a nucleic acid and a pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein, said src protein having any amino acid residue at codon 527 except tyrosine, serine or threonine.

Also envisioned is a pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a viral or non-viral gene transfer vector containing a nucleic acid and a pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein having no kinase activity.

Brief Description of the Drawings

In the drawings forming a portion of this disclosure:

FIG. 1 is a cDNA sequence of chicken c-Src which is the complete coding sequence with the introns deleted as first described by Takeya et al., *Cell*, 32:881-890 (1983). The sequence is accessible through GenBank Accession Number

J00844. The sequence contains 1759 nucleotides with the protein coding portion beginning and ending at the respective nucleotide positions 112 and 1713.

FIG. 2 is the encoded amino acid residue sequence of chicken c-Src of the coding sequence shown in FIG. 1.

5 FIG. 3 is a cDNA sequence of human c-Src which as first described by Braeuninger et al., Proc. Natl. Acad. Sci., USA, 88:10411-10415 (1991). The sequence is accessible through GenBank Accession Number X59932 X71157. The sequence contains 2187 nucleotides with the protein coding portion beginning and ending at the respective nucleotide positions 134 and 1486.

10 FIG. 4 is the encoded amino acid residue sequence of human c-Src of the coding sequence shown in FIG. 3.

FIG. 5 illustrates the activation of endogenous Src by bFGF or VEGF as described in Example 4. The top portion of the figure indicates the results of an in vitro kinase assay with the fold activation of endogenous c-Src by either bFGF and VEGF. The bottom portion of the figure is the kinase assay blot probed with an anti-Src antibody as a loading control for equivalent Src and IgG content.

15 FIG. 6 illustrates the effect of retrovirus-mediated gene expression of c-Src A on angiogenesis in the chick chorioallantoic membrane (CAM) as described in Example 4. Nine-day-old chick CAMs were exposed to RCAS-Src A (active mutated c-Src) or control RCAS-GFP (Green Fluorescent Protein; a fluorescent indicator protein) retroviruses or buffer for 72 h. The level of angiogenesis was quantified as shown in FIG. 6A with representative photomicrographs (4x) in FIG. 6B corresponding to each treatment taken with a stereomicroscope.

20 FIG. 7 illustrates the retroviral expression of c-Src A in activating vascular MAP kinase phosphorylation. FIG. 7A shows tissue extracts of 10 day-old chick CAMs that had been exposed to VEGF or PMA for 30 minutes or infected with c-Src A retrovirus for 48 hours. NT stands for no treatment. Src was immunoprecipitated from equivalent amounts of total protein extract and subjected to an in vitro immune complex kinase assay using a FAK-GST fusion protein as a substrate, electrophoresed and transferred to nitrocellulose. Aliquots of the above whole tissue lysates were also measured for endogenous ERK phosphorylation by immunoblotting with an anti-phospho-ERK antibody. FIG. 7B shows 10 day old CAMs that were infected with either mock RCAS or RCAS containing SRC A.

After two days, CAMs were dissected, cryopreserved in OCT and sectioned at 4 μ m. Sections were immunostained with an anti-phosphorylated ERK antibody (New England Biolabs), washed and detected with a goat anti-rabbit FITC-conjugated secondary antibody. Florescent images were captured on a cooled-
5 CCD camera (Princeton Inst.)

FIG. 8 illustrates the selective requirement for Src activity during VEGF, but not bFGF-induced angiogenesis. Nine day old chick CAMs were exposed to RCAS-Src 251 or control RCAS-GFP retroviruses or buffer for 20 hours and then incubated for an additional 72 hours in the presence or absence of bFGF or
10 VEGF. The level of angiogenesis was quantified FIG. 8A as described above, and representative photomicrographs (6x) were taken with a stereomicroscope as shown in FIG. 8B. FIG. 8C shows a blot probed with an anti-Src antibody to confirm the expression of Src 251 in transfected cells as compared to mock treatments.

FIG. 9 illustrates the results of retroviral delivery of RCAS-Src 251 to human tumors. FIG. 9A is a micrograph that shows human medulloblastoma tumor fragment infected with RCAS-GFP (RCAS-Green Fluorescent Protein) expressing GFP exclusively in the tumor blood vessels (arrowhead) as detected by optical sectioning with a Bio Rad laser confocal scanning microscope (bar=500 μ m).
20 FIG. 9B depicts data from tumors treated with topical application of retrovirus, which were allowed to grow for 3 or 6 days after which they were resected and wet weights determined. Data are expressed as the mean change in tumor weight (from the 50 mg tumor starting weight) +/- SEM of 2 replicates. FIG. 9C depicts in representative micrographs, medulloblastoma tumors surgically removed from
25 the embryos (bar=350 μ m). The lower panels are high magnification views of each tumor showing the vasculature of each tumor in detail (bar=350 μ m). The arrowhead indicates blood vessel disruption in RCAS-Src251-treated tumors.

FIG. 10 is a diagram illustrating a restriction map of the RCASBP (RCAS) vector construct.

30 Detailed Description of the Invention

A. Definitions

Amino Acid Residue: An amino acid formed upon chemical digestion (hydrolysis) of a polypeptide at its peptide linkages. The amino acid residues

described herein are preferably in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired functional property is retained by the polypeptide. NH₂ refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide in keeping with standard polypeptide nomenclature (described in J. Biol. Chem., 243:3552-59 (1969) and adopted at 37 CFR §1.822(b)(2)).

It should be noted that all amino acid residue sequences are represented herein by formulae whose left and right orientation is in the conventional direction of amino-terminus to carboxy-terminus. Furthermore, a dash at the beginning or end of an amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino acid residues.

Polypeptide: refers to a linear array of amino acid residues connected to one another by peptide bonds between the alpha-amino group and carboxy group of contiguous amino acid residues.

Peptide: as used herein refers to a linear array of no more than about 50 amino acid residues connected one to the other as in a polypeptide.

Cyclic peptide: refers to a compound having a ring structure that includes several amide bonds as in a typical peptide. The cyclic peptide can be a "head to tail" homodetic cyclic peptide, or it can contain a heterodetic ring structure in which the ring is closed by disulfide bridges, lactam bridges, thioesters, thioamides, guanidino, and the like linkages.

Protein: refers to a linear array of more than 50 amino acid residues connected one to the other as in a polypeptide.

Fusion protein: refers to a polypeptide containing at least two different polypeptide domains operatively linked by a typical peptide bond ("fused"), where the two domains correspond to peptides not found fused in nature.

Synthetic peptide: refers to a chemically produced chain of amino acid residues linked together by peptide bonds that is free of naturally occurring proteins and fragments thereof.

B. General Considerations

The present invention relates generally to the discovery that angiogenesis is mediated by the tyrosine kinase Src protein, and that angiogenesis can be

modulated by providing either active or inactive Src proteins for potentiating or inhibiting angiogenesis, respectively.

This discovery is important because of the role that angiogenesis, the formation of new blood vessels, plays in a variety of disease processes. Where tissues associated with a disease condition require angiogenesis for tissue growth, it is desirable to inhibit angiogenesis and thereby inhibit the diseased tissue growth. Where injured tissue requires angiogenesis for tissue growth and healing, it is desirable to potentiate or promote angiogenesis and thereby promote tissue healing and growth.

Where the growth of new blood vessels is the cause of, or contributes to, the pathology associated with a diseased tissue, inhibition of angiogenesis reduces the deleterious effects of the disease. By inhibiting angiogenesis, one can intervene in the disease, ameliorate the symptoms, and in some cases cure the disease.

Examples of tissue associated with disease and neovascularization that will benefit from inhibitory modulation of angiogenesis include rheumatoid arthritis, diabetic retinopathy, inflammatory diseases, restenosis, and the like. Where the growth of new blood vessels is required to support growth of a deleterious tissue, inhibition of angiogenesis will reduce the blood supply to the tissue and thereby contribute to reduction in tissue mass based on blood supply requirements.

Examples include growth of tumors where neovascularization is a continual requirement in order that the tumor grow beyond a few millimeters in thickness, and for the establishment of solid tumor metastases.

Where the growth of new blood vessels contributes to healing of tissue, potentiation of angiogenesis assists in healing. Examples include treatment of patients with ischemic limbs in which there is poor circulation in the limbs from diabetes or other conditions. Also contemplated for treatment are patients with chronic wounds that do not heal and therefore could benefit from the increase in vascular cell proliferation and neovascularization.

The methods of the present invention are effective in part because the therapy is highly selective for angiogenesis and not other biological processes.

As described earlier, angiogenesis includes a variety of processes involving neovascularization of a tissue including "sprouting", vasculogenesis, or vessel enlargement, all of which angiogenesis processes are effected by Src protein.

With the exception of traumatic wound healing, corpus luteum formation and embryogenesis, it is believed that the majority of angiogenesis processes are associated with disease processes. Accordingly, the present therapeutic methods are selective for the disease and do not have deleterious side effects.

5 C. Src Proteins

A tyrosine kinase Src protein for use in the present invention can vary depending upon the intended use. The terms "Src protein" or "Src" are used to refer to the various forms of tyrosine kinase Src proteins described herein, either in active or inactive forms.

10 An "active Src protein" refers to any of a variety of forms of src protein which potentiate angiogenesis. Assays to measure potentiation of angiogenesis are described herein, and are not to be construed as limiting. A protein is considered active if the level of angiogenesis is at least 10% greater, preferably 25% greater, and more preferably 50% greater than a control level where no src is added to the
15 assay system. The preferred assay for measuring potentiation is the CAM assay using RCAS viral vector as described in the Examples in which the angiogenic index is calculated by counting branch points. A preferred active Src protein exhibits tyrosine kinase activity as well. Exemplary active Src proteins are described in the Examples, and include Src-A.

20 An "inactive Src protein" refers to any of a variety of forms of Src protein which inhibit angiogenesis. Assays to measure inhibition of angiogenesis are described herein, and are not to be construed as limiting. A protein is considered inactive if the level of angiogenesis is at least 10% lower, preferably 25% lower, and more preferably 50% lower than a control level where no exogenous Src is
25 added to the assay system. The preferred assay for measuring inhibition is the CAM assay using RCAS viral vector as described in the Examples in which the angiogenic index is calculated by counting branch points. A preferred inactive Src protein exhibits reduced tyrosine kinase activity as well. Exemplary inactive Src proteins are described in the Examples, and include Src-251.

30 A Src protein useful in the present invention can be produced in any of a variety of methods including isolation from natural sources including tissue, production by recombinant DNA expression and purification, and the like. Src

protein can also be provided "in situ" by introduction of a gene therapy system to the tissue of interest which then expresses the protein in the tissue.

A gene encoding a Src protein can be prepared by a variety of methods known in the art, and the invention is not to be construed as limiting in this regard. For example, the natural history of Src is well known to include a variety of homologs from mammalian, avian, viral and the like species, and the gene can readily be cloned using cDNA cloning methods from any tissue expressing the protein. A preferred Src for use in the invention is a cellular protein, such as the mammalian or avian homologs designated c-Src. Particularly preferred is human c-Src.

10 D. Recombinant DNA Molecules and Expression Systems for Expression of a Src Protein

The invention describes several nucleotide sequences of particular use in the present invention. These sequences include sequences which encode a Src protein useful in the invention, and various DNA segments, recombinant DNA (rDNA) molecules and vectors constructed for expression of Src protein.

DNA molecules (segments) of this invention therefore can comprise sequences which encode whole structural genes, fragments of structural genes, and transcription units as described further herein.

A preferred DNA segment is a nucleotide sequence which encodes a Src protein as defined herein, or biologically active fragment thereof,

The amino acid residue sequence and nucleotide sequence of a preferred c-Src is described in the Examples.

A preferred DNA segment codes for an amino acid residue sequence substantially the same as, and preferably consisting essentially of, an amino acid residue sequence or portions thereof corresponding to a Src protein described herein. Representative and preferred DNA segments are further described in the Examples.

The amino acid residue sequence of a protein or polypeptide is directly related via the genetic code to the deoxyribonucleic acid (DNA) sequence of the structural gene that codes for the protein. Thus, a structural gene or DNA segment can be defined in terms of the amino acid residue sequence, i.e., protein or polypeptide, for which it codes.

An important and well known feature of the genetic code is its redundancy. That is, for most of the amino acids used to make proteins, more than one coding nucleotide triplet (codon) can code for or designate a particular amino acid residue. Therefore, a number of different nucleotide sequences may code for a particular amino acid residue sequence. Such nucleotide sequences are considered functionally equivalent since they can result in the production of the same amino acid residue sequence in all organisms. Occasionally, a methylated variant of a purine or pyrimidine may be incorporated into a given nucleotide sequence. However, such methylations do not affect the coding relationship in any way.

A nucleic acid is any polynucleotide or nucleic acid fragment, whether it be a polyribonucleotide or polynucleotide, i.e., RNA or DNA, or analogs thereof. In preferred embodiments, a nucleic acid molecule is in the form of a segment of duplex DNA, i.e., a DNA segment, although for certain molecular biological methodologies, single-stranded DNA or RNA is preferred.

DNA segments are produced by a number of means including chemical synthesis methods and recombinant approaches, preferably by cloning or by polymerase chain reaction (PCR). DNA segments that encode portions of a Src protein can easily be synthesized by chemical techniques, for example, the phosphotriester method of Matteucci et al, *J. Am. Chem. Soc.*, 103:3185-3191, 1981, or using automated synthesis methods. In addition, larger DNA segments can readily be prepared by well known methods, such as synthesis of a group of oligonucleotides that define the DNA segment, followed by hybridization and ligation of oligonucleotides to build the complete segment. Alternative methods include isolation of a preferred DNA segment by PCR with a pair of oligonucleotide primers used on a cDNA library believed to contain members which encode a Src protein.

Of course, through chemical synthesis, any desired modifications can be made simply by substituting the appropriate bases for those encoding the native amino acid residue sequence. This method is well known, and can be readily applied to the production of the various different "modified" Src proteins described herein.

Furthermore, DNA segments consisting essentially of structural genes encoding a Src protein can be subsequently modified, as by site-directed or random mutagenesis, to introduce any desired substitutions.

1. Cloning a Src Gene

A Src gene can be cloned from a suitable source of genomic DNA or messenger RNA (mRNA) by a variety of biochemical methods. Cloning of these genes can be conducted according to the general methods described in the

5 Examples and as known in the art.

Sources of nucleic acids for cloning a Src gene suitable for use in the methods of this invention can include genomic DNA or messenger RNA (mRNA) in the form of a cDNA library, from a tissue believed to express these proteins. A preferred tissue is human lung tissue, although any other suitable tissue may be

10 used.

A preferred cloning method involves the preparation of a cDNA library using standard methods, and isolating the Src-encoding nucleotide sequence by PCR amplification using paired oligonucleotide primers based on the nucleotide sequences described herein. Alternatively, the desired cDNA clones can be

15 identified and isolated from a cDNA or genomic library by conventional nucleic acid hybridization methods using a hybridization probe based on the nucleic acid sequences described herein. Other methods of isolating and cloning suitable src encoding nucleic acids are readily apparent to one skilled in the art.

2. Expression Vectors

20 A recombinant DNA molecule (rDNA) containing a DNA segment encoding a Src protein can be produced as described herein. In particular, an expressible rDNA can be produced by operatively (in frame, expressibly) linking a vector to a src encoding DNA segment. Thus, a recombinant DNA molecule is a hybrid DNA molecule comprising at least two nucleic acids of a nucleotide sequences not

25 normally found together in nature.

The choice of vector to which the DNA segment is operatively linked depends directly, as is well known in the art, on the functional properties desired, e.g., protein expression, and the host cell to be transformed. A vector suitable for use in practicing the present invention is at least capable of directing the replication,

30 and preferably also expression, of a structural gene included in the vector DNA segments to which it is operatively linked.

Both prokaryotic and eukaryotic expression vectors are familiar to one of ordinary skill in the art of vector construction, and are described by Ausubel, et

al., in Current Protocols in Molecular Biology, Wiley and Sons, New York (1993) and by Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, (1989). These references also describe many of the general recombinant DNA methods referred to herein.

5 In one embodiment, a suitable vector includes a procaryotic replicon, i.e., a DNA sequence having the ability to direct autonomous replication and maintenance of the recombinant DNA molecule extrachromosomally in a procaryotic host cell, such as a bacterial host cell, transformed therewith. Such replicons are well known in the art. In addition, those embodiments that include a procaryotic
10 replicon also include a gene whose expression confers drug resistance to a bacterial host transformed therewith. Typical bacterial drug resistance genes are those that confer resistance to ampicillin or tetracycline.

Those vectors that include a procaryotic replicon can also include a procaryotic promoter capable of directing the expression (transcription and
15 translation) of a structural gene in a bacterial host cell, such as E. coli, transformed therewith. A promoter is an expression control element formed by a DNA sequence that permits binding of RNA polymerase and transcription to occur. Promoter sequences compatible with bacterial hosts are typically provided in plasmid vectors containing convenient restriction sites for insertion of a DNA
20 segment of the present invention. Typical of such vector plasmids are pUC8, pUC9, pBR322 and pBR329 available from Biorad Laboratories, (Richmond, CA), pRSET available from Invitrogen (San Diego, CA) and pPL and pKK223 available from Pharmacia, Piscataway, N.J.

Expression vectors compatible with eukaryotic cells, preferably those
25 compatible with vertebrate cells, can also be used to form the recombinant DNA molecules of the present invention. Eukaryotic cell expression vectors are well known in the art and are available from several commercial sources. Typically, such vectors are provided containing convenient restriction sites for insertion of the desired DNA segment. Typical of such vectors are pSVL and pKSV-10
30 (Pharmacia), pBPV-1/pML2d (International Biotechnologies, Inc.), pTDT1 (ATCC, #31255), pRc/CMV (Invitrogen, Inc.), the preferred vector described in the Examples, and the like eukaryotic expression vectors.

A particularly preferred system for gene expression in the context of this invention includes a gene delivery component, that is, the ability to deliver the gene to the tissue of interest. Suitable vectors are "infectious" vectors such as recombinant DNA viruses, adenovirus or retrovirus vectors which are engineered
5 to express the desired protein and have features which allow infection of preselected target tissues. Particularly preferred is the replication competent avian sarcoma virus (RCAS) described herein.

Mammalian cell systems that utilize recombinant viruses or viral elements to direct expression may be engineered. For example, when using adenovirus
10 expression vectors, the coding sequence of a polypeptide may be ligated to an adenovirus transcription/ translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted into the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a
15 recombinant virus that is viable and capable of expressing the polypeptide in infected hosts (e.g., see Logan et al., Proc. Natl. Acad. Sci., USA, 81:3655-3659 (1984)). Alternatively, the vaccinia virus 7.5K promoter may be used (e.g., see, Mackett et al., Proc. Natl. Acad. Sci., USA, 79:7415-7419 (1982); Mackett et al., J. Virol., 49:857-864 (1984); Panicali et al., Proc. Natl. Acad. Sci., USA,
20 79:4927-4931 (1982)). Of particular interest are vectors based on bovine papilloma virus which have the ability to replicate as extrachromosomal elements (Sarver et al., Mol. Cell. Biol., 1:486 (1981)). Shortly after entry of this DNA into target cells, the plasmid replicates to about 100 to 200 copies per cell. Transcription of the inserted cDNA does not require integration of the plasmid
25 into the host's chromosome, thereby yielding a high level of expression. These vectors can be used for stable expression by including a selectable marker in the plasmid, such as the neo gene. Alternatively, the retroviral genome can be modified for use as a vector capable of introducing and directing the expression of the polypeptide-encoding nucleotide sequence in host cells (Cone et al., Proc.
30 Natl. Acad. Sci., USA, 81:6349-6353 (1984)). High level expression may also be achieved using inducible promoters, including, but not limited to, the metallothioneine IIA promoter and heat shock promoters.

Recently, long-term survival of cytomegalovirus (CMV) promoter versus Rous sarcoma virus (RSV) promoter-driven thymidine kinase (TK) gene therapy in nude mice bearing human ovarian cancer has been studied. Cell killing efficacy of adenovirus-mediated CMV promoter-driven herpes simplex virus TK gene therapy was found to be 2 to 10 time more effective than RSV driven therapy. (Tong et al., 1999, Hybridoma 18(1):93-97). The design of chimeric promoters for gene therapy applications, which call for low level expression followed by inducible high-level expression has also been described. (Suzuki et al., 1996, Human Gene Therapy 7:1883-1893).

For long-term, high-yield production of recombinant proteins, stable expression is preferred. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with a cDNA controlled by appropriate expression control elements (e.g., promoter and enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. As mentioned above, the selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines.

For example, following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., Cell, 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska et al, Proc. Natl. Acad. Sci., USA, 48:2026 (1962)), and adenine phosphoribosyltransferase (Lowy et al., Cell, 22:817 (1980)) genes, which can be employed in tk-, hgp^rt- or ap^rt- cells respectively. Also, antimetabolite resistance-conferring genes can be used as the basis of selection; for example, the genes for dhfr, which confers resistance to methotrexate (Wigler et al., Proc. Natl. Acad. Sci., USA, 77:3567 (1980); O'Hare et al., Proc. Natl. Acad. Sci., USA, 78:1527 (1981); gpt, which confers resistance to mycophenolic acid (Mulligan et al, Proc. Natl. Acad. Sci., USA, 78:2072, (1981)); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al, J. Mol. Biol., 150:1 (1981)); and hyg^r, which confers resistance to hygromycin (Santerre et al, Gene,

30:147 (1984)). Recently, additional selectable genes have been described, namely trpB, which allows cells to utilize indole in place of tryptophan; hisD, which allows cells to utilize histinol in place of histidine (Hartman et al, Proc. Natl. Acad. Sci., USA, 85:804 (1988)); and ODC (ornithine decarboxylase) which
5 confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue L., In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory ed., (1987)).

The principal vectors contemplated for human gene therapy are derived from retroviral origin. (Wilson, 1997, Clin. Exp. Immunol. 107(Sup. 1):31-32; Bank et al., 1996, Bioessays 18(12):999-1007; Robbins et al., 1998, Pharmacol. Ther. 80(1):35-47). The therapeutic potential of gene transfer and antisense therapy has stimulated the development of many vector systems for treating a variety of tissues. (vasculature, Stephan et al., 1997, Fundam. Clin. Pharmacol. 11(2):97-
10 110; Feldman et al., 1997, Cardiovasc. Res. 35(3):391-404; Vassalli et al., 1997, Cardiovasc. Res. 35(3):459-69; Baek et al., 1998, Circ. Res. 82(3):295-305; kidney, Lien et al., 1997, Kidney Int. Suppl. 61:S85-8; liver, Ferry et al., 1998, Hum Gene Ther. 9(14):1975-81; muscle, Marshall et al., 1998, Curr. Opin. Genet. Dev. 8(3):360-5). In addition to these tissues, a critical target for human gene
15 therapy is cancer, either the tumor itself, or associated tissues. (Runnebaum, 1997, Anticancer Res. 17(4B):2887-90; Spear et al., 1998, J. Neurovirol. 4(2):133-47).

Specific examples of viral gene therapy vector systems readily adaptable for use in the methods of the present invention are briefly described below. Retroviral gene delivery has been recently reviewed by Federspiel and Hughes (1998,
25 Methods in Cell Biol. 52:179-214) which describes in particular, the avian leukemia virus (ALV) retrovirus family (Federspiel et al., Proc. Natl. Acad. Sci., USA, 93:4931 (1996); Federspiel et al., Proc. Natl. Acad. Sci., USA, 91:11241 (1994)). Retroviral vectors, including ALV and murine leukemia virus (MLV) are further described by Svoboda (1998, Gene 206:153-163).

30 Modified retroviral/adenoviral expression systems can be readily adapted for practice of the methods of the present invention. For example, murine leukemia virus (MLV) systems are reviewed by Karavanas et al., 1998, Crit. Rev. in Oncology/Hematology 28:7-30. Adenovirus expression systems are reviewed by

Von Seggern and Nemerow in Gene Expression Systems (ed. Fernandez & Hoeffler, Academic Press, San Diego, CA, 1999, chapter 5, pages 112-157).

- Protein expression systems have been demonstrated to have effective use both in vivo and in vitro. For example, efficient gene transfer to human squamous cell carcinomas by a herpes simplex virus (HSV) type 1 amplicon vector has been described. (Carew et al., 1998, Am. J. Surg. 176:404-408). Herpes simplex virus has been used for gene transfer to the nervous system. (Goins et al., 1997, J. Neurovirol. 3 (Sup. 1):S80-8). Targeted suicide vectors using HSV-TK has been tested on solid tumors. (Smiley et al., 1997, Hum. Gene Ther. 8(8):965-77).
- Herpes simplex virus type 1 vector has been used for cancer gene therapy on colon carcinoma cells. (Yoon et al., 1998, Ann. Surg. 228(3):366-74). Hybrid vectors have been developed to extend the length of time of transfection, including HSV/AAV (adeno-associated virus) hybrids for treating hepatocytes. (Fraefel et al., 1997, Mol. Med. 3(12):813-825).
- Vaccinia virus has been developed for human gene therapy because of its large genome. (Peplinski et al., 1998, Surg. Oncol. Clin. N. Am. 7(3):575-88). Thymidine kinase-deleted vaccinia virus expressing purine nucleoside pyrophosphorylase has been described for use as a tumor directed gene therapy vector. (Puhlman et al., 1999, Human Gene Therapy 10:649-657).
- Adeno-associated virus 2 (AAV) has been described for use in human gene therapy, however AAV requires a helper virus (such as adenovirus or herpes virus) for optimal replication and packaging in mammalian cells. (Snoeck et al., 1997, Exp. Nephrol. 5(6):514-20; Rabinowitz et al., 1998, Curr. Opin. Biotechnol. 9(5):470-5). However, in vitro packaging of an infectious recombinant AAV has been described, making this system much more promising. (Ding et al., 1997, Gene Therapy 4:1167-1172). It has been shown that the AAV mediated transfer of ecotropic retrovirus receptor cDNA allows ecotropic retroviral transduction of established and primary human cells. (Qing et al., 1997, J. Virology 71(7):5663-5667). Cancer gene therapy using an AAV vector expressing human wild-type p53 has been demonstrated. (Qazilbash et al., 1997, Gene Therapy 4:675-682). Gene transfer into vascular cells using AAV vectors has also been shown. (Maeda et al., 1997, Cardiovascular Res. 35:514-521). AAV has been demonstrated as a suitable vector for liver directed gene therapy. (Xiao et al., 1998, J. Virol. 72(12):10222-

6). AAV vectors have been demonstrated for use in gene therapy of brain tissues and the central nervous system. (Chamberlin et al., 1998, Brain Res. 793(1-2):169-75; During et al., 1998, Gene Therapy 5(6):820-7). AAV vectors have also been compared with adenovirus vectors (AdV) for gene therapy of the lung and transfer to human cystic fibrosis epithelial cells. (Teramoto et al., 1998, J. Virol. 72(11):8904-12).

Chimeric AdV/retroviral gene therapy vector systems which incorporate the useful qualities of each virus to create a nonintegrative AdV that is rendered functionally integrative via the intermediate generation of a retroviral producer cell. (Feng et al., 1997, Nat. Biotechnology 15(9):866-70; Bilbao et al., 1997, FASEB J 11(8):624-34). This powerful new generation of gene therapy vector has been adapted for targeted cancer gene therapy. (Bilbao et al., 1998, Adv. Exp. Med. Biol. 451:365-74). Single injection of AdV expressing p53 inhibited growth of subcutaneous tumor nodules of human prostate cancer cells. (Asgari et al., 1997, Int. J. Cancer 71(3):377-82). AdV mediated gene transfer of wild-type p53 in patients with advanced non-small cell lung cancer has been described. (Schuler et al., 1998, Human Gene Therapy 9:2075-2082). This same cancer has been the subject of p53 gene replacement therapy mediated by AdV vectors. (Roth et al., 1998, Semin. Oncol. 25(3 Suppl 8):33-7). AdV mediated gene transfer of p53 inhibits endothelial cell differentiation and angiogenesis in vivo. (Riccioni et al., 1998, Gene Ther. 5(6):747-54). Adenovirus-mediated expression of melanoma antigen gp75 as immunotherapy for metastatic melanoma has also been described. (Hirschowitz et al., 1998, Gene Therapy 5:975-983). AdV facilitates infection of human cells with ecotropic retrovirus and increases efficiency of retroviral infection. (Scott-Taylor, et al., 1998, Gene Ther. 5(5):621-9). AdV vectors have been used for gene transfer to vascular smooth muscle cells (Li et al., 1997, Chin. Med. J.(Engl) 110(12):950-4), squamous cell carcinoma cells (Goebel et al., 1998, Otolaryngol Head Neck Surg 119(4):331-6), esophageal cancer cells (Senmaru et al., 1998, Int J. Cancer 78(3):366-71), mesangial cells (Nahman et al., 1998, J. Investig. Med. 46(5):204-9), glial cells (Chen et al., 1998, Cancer Res. 58(16):3504-7), and to the joints of animals (Ikeda et al., 1998, J. Rheumatol. 25(9):1666-73). More recently, catheter-based pericardial gene transfer mediated by AcV vectors has been demonstrated. (March et al., 1999, Clin. Cardiol. 22(1

Suppl 1):123-9). Manipulation of the AdV system with the proper controlling genetic elements allows for the AdV-mediated regulable target gene expression in vivo. (Burcin et al., 1999, PNAS (USA) 96(2):355-60).

Alphavirus vectors have been developed for human gene therapy applications, with packaging cell lines suitable for transformation with expression cassettes suitable for use with Sindbis virus and Semliki Forest virus-derived vectors. (Polo et al., 1999, Proc. Natl. Acad. Sci., USA, 96:4598-4603). Noncytopathic flavivirus replicon RNA-based systems have also been developed. (Varnavski et al., 1999, Virology 255(2):366-75). Suicide HSV-TK gene containing sinbis virus vectors have been used for cell-specific targeting into tumor cells. (Iijima et al., 1998, Int. J. Cancer 80(1):110-8).

Retroviral vectors based on human foamy virus (HFV) also show promise as gene therapy vectors. (Trobridge et al., 1998, Human Gene Therapy 9:2517-2525). Foamy virus vectors have been designed for suicide gene therapy. (Nestler et al., 1997, Gene Ther. 4(11):1270-7). Recombinant murine cytomegalovirus and promoter systems have also been used as vectors for high level expression. (Manning et al., 1998, J. Virol. Meth. 73(1):31-9; Tong et al., 1998, Hybridoma 18(1):93-7).

Gene delivery into non-dividing cells has been made feasible by the generation of Sendai virus based vectors. (Nakanishi et al., 1998, J. Controlled Release 54(1):61-8).

In other efforts to enable the transformation of non-dividing somatic cells, lentiviral vectors have been explored. Gene therapy of cystic fibrosis using a replication-defective human immunodeficiency virus (HIV) based vector has been described. (Goldman et al., 1997, Human Gene Therapy 8:2261-2268). Sustained expression of genes delivered into liver and muscle by lentiviral vectors has also been shown. (Kafri et al., 1997, Nat. Genet. 17(3):314-7). However, safety concerns are predominant, and improved vector development is proceeding rapidly. (Kim et al., 1998, J. Virol. 72(2):994-1004). Examination of the HIV LTR and Tat yield important information about the organization of the genome for developing vectors. (Sadaie et al., 1998, J. Med. Virol. 54(2):118-28). Thus the genetic requirements for an effective HIV based vector are now better understood. (Gasmi et al., 1999, J. Virol. 73(3):1828-34). Self inactivating vectors, or

conditional packaging cell lines have been described. (for example Zuffery et al., 1998, J. Virol. 72(12):9873-80; Miyoshi et al., 1998, J. Virol. 72(10):8150-7; Dull et al., 1998, J. Virol. 72(11):8463-71; and Kaul et al., 1998, Virology 249(1):167-74). Efficient transduction of human lymphocytes and CD34+ cells by
5 HIV vectors has been shown. (Douglas et al., 1999, Hum. Gene Ther. 10(6):935-45; Miyoshi et al., 1999, Science 283(5402):682-6). Efficient transduction of nondividing human cells by feline immunodeficiency virus (FIV) lentiviral vectors has been described, which minimizes safety concerns with using HIV based vectors. (Poeschla et al., 1998, Nature Medicine 4(3):354-357). Productive
10 infection of human blood mononuclear cells by FIV vectors has been shown. (Johnston et al., 1999, J. Virol. 73(3):2491-8).

While many viral vectors are difficult to handle, and capacity for inserted DNA limited, these limitations and disadvantages have been addressed. For example, in addition to simplified viral packaging cell lines, Mini-viral vectors,
15 derived from human herpes virus, herpes simplex virus type 1 (HSV-1), and Epstein-Barr virus (EBV), have been developed to simplify manipulation of genetic material and generation of viral vectors. (Wang et al., 1996, J. Virology 70(12):8422-8430). Adaptor plasmids have been previously shown to simplify insertion of foreign DNA into helper-independent Retroviral vectors. (1987, J.
20 Virology 61(10):3004-3012).

Viral vectors are not the only means for effecting gene therapy, as several non-viral vectors have also been described. A targeted non-viral gene delivery vector based on the use of Epidermal Growth Factor/DNA polyplex (EGF/DNA) has been shown to result in efficient and specific gene delivery. (Cristiano, 1998,
25 Anticancer Res. 18:3241-3246). Gene therapy of the vasculature and CNS have been demonstrated using cationic liposomes. (Yang et al., 1997, J. Neurotrauma 14(5):281-97). Transient gene therapy of pancreatitis has also been accomplished using cationic liposomes. (Denham et al., 1998, Ann. Surg. 227(6):812-20). A chitosan-based vector/DNA complexes for gene delivery have been shown to be
30 effective. (Erbacher et al., 1998, Pharm. Res. 15(9):1332-9). A non-viral DNA delivery vector based on a terplex system has been described. (Kim et al., 1998, 53(1-3):175-82). Virus particle coated liposome complexes have also been used to

effect gene transfer. (Hirai et al., 1997, Biochem. Biophys. Res. Commun. 241(1):112-8).

Cancer gene therapy by direct tumor injections of nonviral T7 vector encoding a thymidine kinase gene has been demonstrated. (Chen et al., 1998, Human Gene Therapy 9:729-736). Plasmid DNA preparation is important for direct injection gene transfer. (Horn et al., 1995, Hum. Gene Ther. 6(5):656-73). Modified plasmid vectors have been adapted specifically for direct injection. (Hartikka et al., 1996, Hum. Gene Ther. 7(10):1205-17).

Thus, a wide variety of gene transfer/gene therapy vectors and constructs are known in the art. These vectors are readily adapted for use in the methods of the present invention. By the appropriate manipulation using recombinant DNA/molecular biology techniques to insert an operatively linked src (either active or inactive) into the selected expression/delivery vector, many equivalent vectors for the practice of the present invention can be generated.

15 E. Methods For Modulation of Angiogenesis

The invention provides for a method for the modulation of angiogenesis in a tissue associated with a disease process or condition, and thereby effect events in the tissue which depend upon angiogenesis. Generally, the method comprises administering to the tissue associated with a disease process or condition, a composition comprising an angiogenesis-modulating amount of a Src protein or nucleic acid vector expressing active or inactive Src.

As described herein, any of a variety of tissues, or organs comprised of organized tissues, can support angiogenesis in disease conditions including skin, muscle, gut, connective tissue, joints, bones and the like tissue in which blood vessels can invade upon angiogenic stimuli.

The patient treated according to the present invention in its many embodiments is desirably a human patient, although it is to be understood that the principles of the invention indicate that the invention is effective with respect to all mammals, which are intended to be included in the term "patient". In this context, a mammal is understood to include any mammalian species in which treatment of tissue associated with diseases involving angiogenesis is desirable, particularly agricultural and domestic mammalian species.

Thus the method comprises administering to a patient a therapeutically effective amount of a physiologically tolerable composition containing a Src protein or DNA vector for expressing a Src protein in practicing the methods of the invention.

5 The dosage ranges for the administration of a Src protein depend upon the form of the protein, and its potency, as described further herein. The dosage amounts are large enough to produce the desired effect in which angiogenesis and the disease symptoms mediated by angiogenesis are ameliorated. The dosage should not be so large as to cause adverse side effects, however, such as
10 hyperviscosity syndromes, pulmonary edema, congestive heart failure, and the like. Generally, the dosage will vary with the age, condition, sex of the patient, and extent of the disease in the patient, and can be readily determined by one of skill in the art. The dosage can also be adjusted by the individual physician in the event of any complication.

15 A therapeutically effective amount is an amount of Src protein, or nucleic acid encoding for (active or inactive) src protein, sufficient to produce a detectable modulation of angiogenesis in the tissue being treated, ie., an angiogenesis-modulating amount. Modulation of angiogenesis can be measured by CAM assay as described herein, or by other methods known to one skilled in the art.

20 The Src protein or nucleic acid vector expressing the Src protein can be administered parenterally by injection or by gradual infusion over time. Although the tissue to be treated can typically be accessed in the body by systemic administration, and therefore is most often treated by intravenous administration of therapeutic compositions, other tissues and delivery means are contemplated as
25 well. Thus, compositions of the invention can be administered intravenously, intraperitoneally, intramuscularly, subcutaneously, intracavity, transdermally, and can be delivered by peristaltic means.

 The therapeutic compositions containing a Src protein or nucleic acid vector expressing the Src protein can be conventionally administered intravenously, as by
30 injection of a unit dose, for example. The term "unit dose" when used in reference to a therapeutic composition of the present invention refers to physically discrete units suitable as unitary dosage for the subject, each unit containing a

predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required diluent; i.e., carrier, or vehicle.

In one preferred embodiment the reagent is administered in a single dosage intravenously. Localized administration can be accomplished by direct injection or
5 by taking advantage of anatomically isolated compartments, isolating the microcirculation of target organ systems, reperfusion in a circulating system, or catheter based temporary occlusion of target regions of vasculature associated with diseased tissues.

The compositions are administered in a manner compatible with the dosage
10 formulation, and in a therapeutically effective amount. The quantity to be administered and timing depends on the subject to be treated, capacity of the subject's system to utilize the active ingredient, and degree of therapeutic effect desired. Precise amounts of active ingredient required to be administered depend
15 on the judgement of the practitioner and are peculiar to each individual. However, suitable dosage ranges for systemic application are disclosed herein and depend on the route of administration. Suitable regimes for administration are also variable, but are typified by an initial administration followed by repeated doses at one or more hour intervals by a subsequent injection or other administration. Alternatively, continuous intravenous infusion sufficient to maintain concentrations
20 in the blood in the ranges specified for in vivo therapies are contemplated.

1. Inhibition of Angiogenesis

Inhibition of angiogenesis is important in a variety of diseases, referred to as angiogenic diseases. Such diseases include, but are not limited to, inflammatory disorders such as immune and non-immune inflammation, chronic articular
25 rheumatism and psoriasis, disorders associated with inappropriate or inopportune invasion of vessels such as diabetic retinopathy, neovascular glaucoma, restenosis, capillary proliferation in atherosclerotic plaques and osteoporosis, and cancer associated disorders, such as solid tumors, solid tumor metastases, angiofibromas, retrolental fibroplasia, hemangiomas, Kaposi sarcoma and the like cancers which
30 require neovascularization to support tumor growth.

Thus, methods which inhibit angiogenesis in a tissue associated with a disease condition ameliorates symptoms of the disease and, depending upon the disease, can contribute to cure of the disease. In one embodiment, the invention

contemplates inhibition of angiogenesis, per se, in a tissue associated with a disease condition. The extent of angiogenesis in a tissue, and therefore the extent of inhibition achieved by the present methods, can be evaluated by a variety of methods.

5 Thus, in one related embodiment, a tissue to be treated is an inflamed tissue and the angiogenesis to be inhibited is inflamed tissue angiogenesis where there is neovascularization of inflamed tissue. In this class the method contemplates inhibition of angiogenesis in arthritic tissues, such as in a patient with chronic articular rheumatism, in immune or non-immune inflamed tissues, in psoriatic
10 tissue and the like.

 In another related embodiment, a tissue to be treated is a retinal tissue of a patient with a retinal disease such as diabetic retinopathy, macular degeneration or neovascular glaucoma and the angiogenesis to be inhibited is retinal tissue angiogenesis where there is neovascularization of retinal tissue.

15 In an additional related embodiment, a tissue to be treated is a tumor tissue of a patient with a solid tumor, a metastases, a skin cancer, a breast cancer, a hemangioma or angiofibroma and the like cancer, and the angiogenesis to be inhibited is tumor tissue angiogenesis where there is neovascularization of a tumor tissue. Typical solid tumor tissues treatable by the present methods include lung,
20 pancreas, breast, colon, laryngeal, ovarian, and the like tissues. Inhibition of tumor tissue angiogenesis is a particularly preferred embodiment because of the important role neovascularization plays in tumor growth. In the absence of neovascularization of tumor tissue, the tumor tissue does not obtain the required nutrients, slows in growth, ceases additional growth, regresses and ultimately
25 becomes necrotic resulting in killing of the tumor.

 Stated in other words, the present invention provides for a method of inhibiting tumor neovascularization by inhibiting tumor angiogenesis according to the present methods. Similarly, the invention provides a method of inhibiting tumor growth by practicing the angiogenesis-inhibiting methods.

30 The methods are also particularly effective against the formation of metastases because (1) their formation requires vascularization of a primary tumor so that the metastatic cancer cells can exit the primary tumor and (2) their establishment in a secondary site requires neovascularization to support growth of the metastases.

In a related embodiment, the invention contemplates the practice of the method in conjunction with other therapies such as conventional chemotherapy directed against solid tumors and for control of establishment of metastases. The administration of angiogenesis inhibitor is typically conducted during or after chemotherapy, although it is preferably to inhibit angiogenesis after a regimen of chemotherapy at times where the tumor tissue will be responding to the toxic assault by inducing angiogenesis to recover by the provision of a blood supply and nutrients to the tumor tissue. In addition, it is preferred to administer the angiogenesis inhibition methods after surgery where solid tumors have been removed as a prophylaxis against metastases.

Insofar as the present methods apply to inhibition of tumor neovascularization, the methods can also apply to inhibition of tumor tissue growth, to inhibition of tumor metastases formation, and to regression of established tumors.

Restenosis is a process of smooth muscle cell (SMC) migration and proliferation into the tissue at the site of percutaneous transluminal coronary angioplasty which hampers the success of angioplasty. The migration and proliferation of SMC's during restenosis can be considered a process of angiogenesis which is inhibited by the present methods. Therefore, the invention also contemplates inhibition of restenosis by inhibiting angiogenesis according to the present methods in a patient following angioplasty procedures. For inhibition of restenosis, the inactivated tyrosine kinase is typically administered after the angioplasty procedure because the coronary vessel wall is at risk of restenosis, typically for from about 2 to about 28 days, and more typically for about the first 14 days following the procedure.

The present method for inhibiting angiogenesis in a tissue associated with a disease condition, and therefore for also practicing the methods for treatment of angiogenesis-related diseases, comprises contacting a tissue in which angiogenesis is occurring, or is at risk for occurring, with a composition comprising a therapeutically effective amount of an inactivated Src protein or vector expressing the protein.

Inhibition of angiogenesis and tumor regression occurs as early as 7 days after the initial contacting with the therapeutic composition. Additional or prolonged

exposure to inactive Src protein is preferable for 7 days to 6 weeks, preferably about 14 to 28 days.

2. Potentialiation of Angiogenesis

In cases where it is desirable to promote or potentiate angiogenesis, administration of an active Src protein to the tissue is useful. The routes and timing of administration are comparable to the methods described hereinabove for inhibition.

F. Therapeutic Compositions

The present invention contemplates therapeutic compositions useful for practicing the therapeutic methods described herein. Therapeutic compositions of the present invention contain a physiologically tolerable carrier together with a Src protein or vector capable of expressing a Src protein as described herein, dissolved or dispersed therein as an active ingredient. In a preferred embodiment, the therapeutic composition is not immunogenic when administered to a mammal or human patient for therapeutic purposes.

As used herein, the terms "pharmaceutically acceptable", "physiologically tolerable" and grammatical variations thereof, as they refer to compositions, carriers, diluents and reagents, are used interchangeably and represent that the materials are capable of administration to or upon a mammal without the production of undesirable physiological effects such as nausea, dizziness, gastric upset and the like.

The preparation of a pharmacological composition that contains active ingredients dissolved or dispersed therein is well understood in the art and need not be limited based on formulation. Typically such compositions are prepared as injectable either as liquid solutions or suspensions, however, solid forms suitable for solution, or suspensions, in liquid prior to use can also be prepared. The preparation can also be emulsified or presented as a liposome composition.

The active ingredient can be mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient and in amounts suitable for use in the therapeutic methods described herein. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol or the like and combinations thereof. In addition, if desired, the composition can contain minor amounts of

auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like which enhance the effectiveness of the active ingredient.

The therapeutic composition of the present invention can include pharmaceutically acceptable salts of any salt-forming components therein.

- 5 Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide) that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, tartaric, mandelic and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, 10 ammonium, calcium or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine and the like.

- Physiologically tolerable carriers for the active ingredients are well known in the art. Exemplary of liquid carriers are sterile aqueous solutions that contain no 15 materials in addition to the active ingredients and water, or contain a buffer such as sodium phosphate at physiological pH value, physiological saline or both, such as phosphate-buffered saline. Still further, aqueous carriers can contain more than one buffer salt, as well as salts such as sodium and potassium chlorides, dextrose, polyethylene glycol and other solutes.

- 20 Liquid compositions can also contain liquid phases in addition to and to the exclusion of water. Exemplary of such additional liquid phases are glycerin, vegetable oils such as cottonseed oil, and water-oil emulsions.

- A therapeutic composition contains an angiogenesis-modulating amount of an Src protein of the present invention, or sufficient recombinant DNA expression 25 vector to express an effective amount of Src protein, typically formulated to contain an amount of at least 0.1 weight percent of Src protein per weight of total therapeutic composition. A weight percent is a ratio by weight of Src protein to total composition. Thus, for example, 0.1 weight percent is 0.1 grams of Src protein per 100 grams of total composition. For DNA expression vectors, the 30 amount administered depends on the properties of the expression vector, the tissue to be treated, and the like considerations.

G. Article of Manufacture

The invention also contemplates an article of manufacture which is a labelled container for providing a Src protein of the invention. An article of manufacture comprises packaging material provided with appropriate labeling for the disease
5 condition to be treated and a pharmaceutical agent contained within the packaging material.

The pharmaceutical agent in an article of manufacture is any of the compositions of the present invention suitable for providing a Src protein and formulated into a pharmaceutically acceptable form as described herein according
10 to the disclosed indications. Thus, the composition can comprise a Src protein or a DNA molecule which is capable of expressing a Src protein. The article of manufacture contains an amount of pharmaceutical agent sufficient for use in treating a condition indicated herein, either in unit or multiple dosages.

The packaging material comprises a label which indicates the use of the
15 pharmaceutical agent contained therein, e.g., for treating conditions assisted by the inhibition or potentiation of angiogenesis, and the like conditions disclosed herein. The label can further include instructions for use and related information as may be required for marketing. The packaging material can include container(s) for storage of the pharmaceutical agent.

20 As used herein, the term packaging material refers to a material such as glass, plastic, paper, foil, and the like capable of holding within fixed means a pharmaceutical agent. Thus, for example, the packaging material can be plastic or glass vials, laminated envelopes and the like containers used to contain a pharmaceutical composition including the pharmaceutical agent.

25 In preferred embodiments, the packaging material includes a label that is a tangible expression describing the contents of the article of manufacture and the use of the pharmaceutical agent contained therein.

Examples

30 The following examples relating to this invention are illustrative and should not, of course, be construed as specifically limiting the invention. Moreover, such variations of the invention, now known or later developed, which would be within

the purview of one skilled in the art are to be considered to fall within the scope of the present invention hereinafter claimed.

1. Preparation of c-Src Expression Constructs

5 For preparing the expression constructs useful in modulating angiogenesis by the methods of the present invention, c-Src cDNA is manipulated and inserted into an expression construct/vector.

The cDNA sequence encoding for wild-type (i.e., endogenous) chicken c-Src is shown in FIG. 1 (SEQ ID NO.:2) with the encoded amino acid residue sequence shown in FIG. 2 (SEQ ID NO.:3). The encoded protein sequence is translated from the cDNA nucleotide positions 112 to 1713. The nucleic acid sequence corresponding to the nucleic acid sequence of human c-Src cDNA (SEQ ID NO.:4) and encoded amino acid residue (SEQ ID NO.:5) sequences are shown respectively in FIGs. 3 and 4. For the human protein sequence, the coding sequence begins at nucleotide position 134 to 1486 of the cDNA.

15 Wild-type as well as a number of mutated c-Src cDNAs were prepared. Mutated c-Src constructs were prepared by site-directed mutagenesis as described by Kaplan et al., EMBO J., 13:4745-4756 (1994). The mutated c-Src constructs for encoding mutated c-Src proteins for use in the methods of the present invention are described in Kaplan et al., id. Kaplan et al. describe various mutated c-Src constructs and encoded proteins of useful for the practice of this invention. For example, Kaplan et al. depict several products of chicken c-src alleles in their FIG. 1, including SrcA and Src251.

25 Two categories of c-Src function to modulate angiogenesis are described. As previously discussed, one category contains Src molecules that increase angiogenesis and thus are considered to be active proteins. Wild-type Src along with various mutations are shown in the present invention to induce angiogenesis. One preferred mutation of wild type c-src which functions in this context with respect to its ability to induce blood vessel growth and therefore increase tumor weight in vivo is the Src A mutant having a point mutation at amino acid (aa) residue position 527 changing tyrosine 527 to phenylalanine. This site is normally a site for negative regulation by the c-Src kinase, referred to as kinase CSK. When CSK phosphorylates aa527 in the wild-type src, the protein is inactivated.

30

However, in mutated Src A, the regulatory tyrosine converted to phenylalanine thus conferring upon the protein a constitutively (i.e., permanently) active protein not subject to inactivation by phosphorylation.

5 Mutations in src have also been shown to have the opposite modulatory effect on angiogenesis, inhibiting angiogenesis instead of stimulating it. Such mutations are referred to as inactive src mutations. Proteins having mutation that confer this inhibitory activity are also referred to as dominant negative Src proteins in that they inhibit neovascularization, including that resulting from endogenous activity of Src as well as enhanced Src activity resulting from growth factor stimulation.
10 Thus certain mutations of wild type c-src of the present invention can also function as a dominant negative with respect to their ability to block blood vessel growth, and for example, therefore decrease tumor weight in vivo.

Such preferred inhibitory c-Src protein includes the Src 251 in which only the first 251 amino acids of Src are expressed. This construct lacks the entire kinase
15 domain and is therefore referred to as "kinase dead" src protein. A second construct is the Src (K295M) mutation in which the lysine amino acid residue 295 is mutated into a methionine. This point mutation in the kinase domain prevents ATP binding and also blocks kinase-dependent Src functions related to vascular cell and tumor cell signaling and proliferation.

20 For example, for the mutation at residue 527, as long as the resultant mutated amino acid residue is not tyrosine, serine, or threonine, the present invention contemplates that the presence of an alternate amino acid at the desired position will result in a Src protein with a desired active, angiogenesis promoting modulatory activity.

25 With respect to the point mutations, any mutation resulting in the desired inhibitory or stimulatory activity is contemplated for use in this invention. Fusion protein constructs combining the desired src protein (mutation or fragment thereof) with expressed amino acid tags, antigenic epitopes, fluorescent protein, or other such protein or peptides are also contemplated, so long as the desired modulating
30 effect of the src protein is intact.

TABLE I

| <u>Src/Mutation</u> | <u>Src Function</u> | <u>Effect on Angiogenesis</u> |
|---------------------|---------------------|-------------------------------|
| c-Src | + | active stimulates |
| SrcA (T527F) | + | active stimulates |
| 5 Src527(point) | + | active stimulates |
| Src251 | - | inactive inhibits |
| Src (truncate) | - | inactive inhibits |
| Src(K295M) | - | inactive inhibits |
| Src295 (point) | - | inactive inhibits |

10

One preferred expression construct for use in the present invention is the RCASBP(A) construct (SEQ ID NO.:1). This expression vector is based on a series of replication competent avian sarcoma viruses with an enhanced Bryan polymerase (BP) for improved titre, and is specific for the A type envelope glycoprotein expressed on normal avian cells (Reviewed in *Methods in Cell Biology*, 52:179-214 (1997); see also, Hughes et al., 1987, *J. Virol.* 61:3004-3012; Fekete & Cepko, 1993, *Mol. Cellular Biol.* 13(4):2604-2613; Itoh et al., 1996, *Development* 122:291-300; and Stott et al., 1998, *BioTechniques* 24:660-666). The complete sequence of RCASBP(A) (SEQ ID NO.:1) is given in the attached sequence listing, and a restriction map of the construct is depicted as FIG. 10, referred to herein as RCAS.

20

The original Src 251 construct was subcloned by Dr. Pam Schwartzberg, at NIH in Dr. Harold Varmus' laboratory. Briefly, cloning of a src cDNA sequence for expression thereof was accomplished by inserting a linker containing Not I-BstB1-Not I restriction sites into a unique Not I site in the 5' end of Src 251. Src has a unique Cla I site at the 3' end. Digestion of Src 251 with BstB1 and Cla I generated a BstB1-ClaI fragment which was then ligated into the Cla I site on RCASBP(A). A BstB1 overhang allows for ligation with a Cla I overhang that will not be recut with Cla I. The src constructs suitable for use in practicing the present invention are readily obtained in the above vector by first digesting the RCAS vector containing Src 251 with Not I and Cla I (in a DAM+ background) to allow for insertion of a similarly digested Src cDNA. Therefore this initial RCASBP(A) construct containing Src 251 was further used to subclone all other

30

Src constructs as described above and in Kaplan et al. (1994, The EMBO J. 13(20):4745-4756), into RCASBP(A) via a Not I-Cla I fragment generated through the Src 251 construction. To produce the desired c-src mutations in the cDNA, standard site-directed mutagenesis procedures familiar to one of ordinary skill in the art were utilized. PCR primers designed to incorporate the desired mutations were also designed with restriction sites to facilitate subsequent cloning steps. Entire segments of Src encoding nucleic acid sequences are deleted from the nucleic acid constructs through PCR amplification techniques based on the known cDNA sequences of chicken, human and the like homologs of Src and subsequent formation of new constructs.

In one embodiment of the invention, the 3' PCR primer used to amplify src nucleic acids also encodes for an in-frame sequence. Use of this primer adds a 9E10-myc epitope tag to the carboxyl terminus of the subsequent Src construct.

The following amino acids were added after amino acid 251 of Src to generate vector constructs containing the 9E10-myc epitope tag: VDMEQKLIAEEDLN (SEQ ID NO.: 6). Two separate PCRs were carried out for each construct and similar results were obtained. All mutant constructs constructed by PCR were also sequenced by PCR to confirm predicted DNA sequence of clones. Wild-type and mutated Src cDNAs for use in the expression systems of the present invention are also available from Upstate Biotech Laboratories, Lake Placid, NY which sells avian as well as human src, and several kinase dead and activated mutated forms.

Alternative expression vectors for use in the expressing the Src proteins of the present invention also include adenoviral vectors as described in US Patent Numbers 4,797,368, 5,173,414, 5,436,146, 5,589,377, and 5,670,488. Alternative methods for the delivery of the Src modulatory proteins include delivery of the Src cDNA with a non-viral vector system as described in US Patent Number 5,675,954 and delivery of the cDNA itself as naked DNA as described in US Patent Number 5,589,466. Delivery of constructs of this invention is also not limited to topical application of a viral vector as described in the CAM assay system below. For example, viral vector preparations are also injected intravenously for systemic delivery into the vascular bed. These vectors are also targetable to sites of increased neovascularization by localized injection of a tumor, as an example.

In vitro expressed proteins are also contemplated for delivery thereof following expression and purification of the selected Src protein by methods useful for delivery of proteins or polypeptides. One such method includes liposome delivery systems, such as described in US Patent Numbers 4,356,167, 5,580,575, 5,542,935 and 5,643,599. Other vector and protein delivery systems are well known to those of ordinary skill in the art for use in the expression and/or delivery of the Src proteins of the present invention.

2. Characterization of the Untreated Chick Chorioallantoic Membrane (CAM)

A. Preparation of the CAM

Angiogenesis can be induced on the chick chorioallantoic membrane (CAM) after normal embryonic angiogenesis has resulted in the formation of mature blood vessels. Angiogenesis has been shown to be induced in response to specific cytokines or tumor fragments as described by Leibovich et al., Nature, 329:630 (1987) and Ausprunk et al., Am. J. Pathol., 79:597 (1975). CAMs were prepared from chick embryos for subsequent induction of angiogenesis and inhibition thereof. Ten day old chick embryos were obtained from McIntyre Poultry (Lakeside, CA) and incubated at 37°C with 60% humidity. A small hole was made through the shell at the end of the egg directly over the air sac with the use of a small crafts drill (Dremel, Division of Emerson Electric Co. Racine WI). A second hole was drilled on the broad side of the egg in a region devoid of embryonic blood vessels determined previously by candling the egg. Negative pressure was applied to the original hole, which resulted in the CAM (chorioallantoic membrane) pulling away from the shell membrane and creating a false air sac over the CAM. A 1.0 centimeter (cm) x 1.0 cm square window was cut through the shell over the dropped CAM with the use of a small model grinding wheel (Dremel). The small window allowed direct access to the underlying CAM.

The resultant CAM preparation was then either used at 6 days of embryogenesis, a stage marked by active neovascularization, without additional treatment to the CAM reflecting the model used for evaluating effects on embryonic neovascularization or used at 10 days of embryogenesis where angiogenesis has subsided. The latter preparation was thus used in this invention

for inducing renewed angiogenesis in response to cytokine treatment or tumor contact as described below.

3. CAM Angiogenesis Assay

A. Angiogenesis Induced by Growth Factors

5 Angiogenesis has been shown to be induced by cytokines or growth factors.

Angiogenesis was induced by placing a 5 millimeter (mm) X 5 mm Whatman filter disk (Whatman Filter paper No.1) saturated with Hanks Balanced Salt Solution (HBSS, GIBCO, Grand Island, NY) or HBSS containing 2 micrograms/milliliter ($\mu\text{g/ml}$) recombinant basic fibroblast growth factor (bFGF) or vascular endothelial cell growth factor (VEGF) (Genzyme, Cambridge, MA) on the CAM of either a 9 or 10 day chick embryo in a region devoid of blood vessels and the windows were latter sealed with tape. Other concentrations of growth factors are also effective at inducing blood vessel growth. For assays where inhibition of angiogenesis is evaluated with intravenous injections of antagonists, angiogenesis is first induced with 1-2 $\mu\text{g/ml}$ bFGF or VEGF in fibroblast growth medium. Angiogenesis was monitored by photomicroscopy after 72 hours.

B. Embryonic Angiogenesis

The CAM preparation for evaluating the effect of angiogenesis inhibitors on the natural formation of embryonic neovasculature is the 6 day embryonic chick embryo as previously described. At this stage in development, the blood vessels are undergoing de novo growth and thus provides a useful system for assessing angiogenesis modulation by the Src proteins of the present invention. The CAM system is prepared as described above with the exception that the assay is performed at embryonic day 6 rather than at day 9 or 10.

25 4. Modulation of Angiogenesis as Measured in the CAM Assay

To assess the effect of Src proteins on angiogenesis, the following assays were performed on 10 day old chick CAM preparations. Five μg of RCAS constructs prepared as described in Example 1 were transfected into the chicken immortalized fibroblast line, DF-1 (gift of Doug Foster, U. of Minn.). This cell line as well as primary chick embryo fibroblasts were capable of producing virus, however the DF-1 cell line produced higher titres. Viral supernatants were collected from subconfluent DF-1 producer cell lines in serum free CLM media [composition: F-10 media base supplemented with DMSO, folic acid, glutamic acid, and MEM

vitamin solution]. Thirty-five ml of viral supernatant were concentrated by ultracentrifugation at 4°C for 2 hours at 22,000 rpm. These concentrated viral pellets were resuspended in 1/100 the original volume in serum-free CLM media, aliquoted and stored at -80°C. The titre was assessed by serial dilution of a control viral vector having a nucleotide sequence encoding green fluorescent protein (GFP), referred to as RCAS-GFP, infection on primary chick embryo fibroblasts that were incubated for 48-72 hours. The titres of viral stock that were obtained following concentration routinely exceeded 10⁸ I.u./ml. For the CAM assay using the viral stocks, cortisone acetate soaked Whatman filter disks 6 mm in diameter were prepared in 3 mg/ml cortisone acetate for 30 minutes in 95% ethanol. The disks were dried in a laminar flow hood and then soaked on 20 µl of viral stock per disk for 10 minutes. These disks were applied to the CAM of 9 or 10 day chick embryos and sealed with cellophane tape and incubated at 37°C for 18-24 hr. Then either mock PBS or growth factors were added at a concentration of 5 µg/ml to the CAM in a 20 µl volume of the appropriate virus stock as an additional boost of virus to the CAM tissue. After 72 hours, the CAMs were harvested and examined for changes in the angiogenic index as determined by double blind counting of the number of branch points in the CAM underlying the disk. For kinase assays, the tissue underlying the disk was harvested in RIPA, homogenized with a motorized grinder and Src immunoprecipitated from equivalent amounts of total protein and subjected to an *in vitro* kinase assay using a FAK-GST fusion protein as a substrate. For the immunofluorescence studies, CAM tissue underlying the disks were frozen in OCT, a cryopreservative, sectioned at 4 µm, fixed in acetone for 1 minute, incubated in 3% normal goat serum for 1 hour, followed by an incubation in primary rabbit anti-phosphorylated ERK antibody as described previously (Eliceiri et al., *J. Cell Biol.*, 140:1255-1263 (1998), washed in PBS and detected with a fluorescent secondary antibody.

A. Activation of Endogenous Src by bFGF or VEGF

To assess the effects of growth factors on Src activity in modulating angiogenesis, the following assays were performed. Tissue extracts of 10 day old chick CAMs that had been exposed to bFGF or VEGF (2 µg/ml) for 2 hours were lysed. Endogenous Src was immunoprecipitated from equivalent amounts of total protein and subjected to an *in vitro* immune complex kinase assay using a FAK-

GST fusion protein as a substrate, electrophoresed and transferred to nitrocellulose.

The results of the assay are shown in FIG. 5 where the increase in Src activity is evident in the increased density of the gel with either bFGF or VEGF treatment as compared to untreated (mock) samples that are indicative of baseline Src activity in the CAM assay. Both bFGF and VEGF resulted in approximately a 2 fold increase of endogenous Src activity present in the CAM. The above kinase assay blot was also probed with an anti-Src antibody as a loading control for equivalent Src and IgG content.

10 B. Effect of Retrovirus-Mediated Gene Expression of Src A on Angiogenesis in the Chick CAM

The following assay was performed to assess the effect of mutated Src proteins on angiogenesis in the CAM preparation. For the assay, 9 day old chick CAMs were exposed to RCAS-Src A or RCAS-GFP expressing retroviruses or buffer for 72 hour following the protocol described above.

The results of this assay are shown in FIG. 6A where the level of angiogenesis was quantified as described above. Representative photomicrographs (4x) were taken with a stereomicroscope as shown in FIG. 6B. Baseline endogenous Src activity has an angiogenic index of approximately 50. In contrast, CAMs treated with retroviral vector-expressed RCAS-Src A having a point mutation at amino acid residue position 527 from a tyrosine to a phenylalanine resulted in an enhancement (induction) of angiogenesis of an angiogenic index of approximately 90. The enhancement of Src-A mediated angiogenesis is also evident in the photographs shown in FIG. 6B.

25 C. Retroviral Expression of Src A Activates Vascular MAP Kinase Phosphorylation

The effect of Src A as compared to growth factors VEGF and PMA on vascular MAP kinase phosphorylation was also assessed following the assay procedures described above and herein. Tissue extracts of 10 day old chick CAMs exposed to VEGF or PMA (another mitogen at a comparable concentration) for 30 minutes were compared to those infected with Src A-expressing retrovirus for 48 hours. Src was then immunoprecipitated from equivalent amounts of total protein extract and subjected to an in vitro immune complex kinase assay using a

FAK-GST fusion protein as a substrate, electrophoresed and transferred to nitrocellulose.

The results of this assay are shown in FIG. 7A where untreated CAMs (NT) exhibit base-line endogenous Src-mediated vascular MAP kinase phosphorylation.

5 Both VEGF and PMA resulted in an approximate 2 fold increase over baseline. In contrast, Src A enhanced the activity approximately 5 to 10 fold over that seen with untreated samples.

Aliquots of the above whole tissue lysates were also measured for endogenous ERK phosphorylation by immunoblotting with an anti-phospho-ERK antibody as shown in FIG. 7B. For this assessment, 10 day old CAMs were infected with
10 either mock RCAS or RCAS that expresses SRC A. After two days, CAMs were dissected, cryopreserved in OCT and sectioned at 4 μ m. Sections were immunostained with an anti-phosphorylated ERK antibody (New England Biolabs), washed and detected with a goat anti-rabbit FITC-conjugated secondary antibody.
15 Fluorescent images were captured on a cooled-CCD camera (Princeton Inst.). The photomicrographs indicate enhanced immunofluorescence with Src A-treated preparations compared to mock controls.

D. Selective Requirement for Src Activity During VEGF, but Not bFGF-Induced Angiogenesis

20 To assess the effect of Src modulatory activity on growth factor induced angiogenesis, the following assays were performed. Nine day old chick CAMs were exposed to the retroviral vector preparation that expressed the dominant negative Src mutation referred to as Src 251 or Src K295M as previously described. RCAS-Src 251 or control RCAS-GFP retroviruses or buffer CAMS
25 were treated for 20 hours and then incubated for an additional 72 hours in the presence or absence of bFGF or VEGF.

The level of angiogenesis, quantified as described above, is shown in FIG. 8A. Representative photomicrographs (6x), shown in FIG. 8B, were taken with a stereomicroscope. FIG. 8C illustrates a blot probed with an anti-Src antibody to
30 confirm the expression of Src 251 in transfected cells as compared to mock treatments.

The results of the assays described above indicate that both bFGF and VEGF treated CAMS in the presence of RCAS-GFP controls induced angiogenesis over

the Src-mediated baseline angiogenesis seen with mock or untreated CAM preparations. The expressed dominant negative mutant Src 251 was effective at inhibiting VEGF-induced angiogenesis back to baseline levels while not effective at inhibiting bFGF-mediated angiogenesis. The photomicrographs shown in FIG. 8B pictorially confirm the data shown in FIG. 8A. Thus, retrovirally expressed Src 251 is an effective angiogenesis inhibitor, when angiogenesis is induced with VEGF.

Applications of the Src proteins of this invention with other angiogenesis models as described in the Examples below are contemplated in the present invention.

5. Regression of Tumor Tissue Growth With Src Modulators as Measured by In Vivo Rabbit Eye Model Assay

The effect of Src modulators on growth factor-induced angiogenesis can be observed in naturally transparent structures as exemplified by the cornea of the eye. New blood vessels grow from the rim of the cornea, which has a rich blood supply, toward the center of the cornea, which normally does not have a blood supply. Stimulators of angiogenesis, such as bFGF, when applied to the cornea induce the growth of new blood vessels from the rim of the cornea. Antagonists of angiogenesis, applied to the cornea, inhibit the growth of new blood vessels from the rim of the cornea. Thus, the cornea undergoes angiogenesis through an invasion of endothelial cells from the rim of the cornea into the tough collagen-packed corneal tissue which is easily visible. The rabbit eye model assay therefore provides an *in vivo* model for the direct observation of stimulation and inhibition of angiogenesis following the implantation of compounds directly into the cornea of the eye.

A. In Vivo Rabbit Eye Model Assay

1) Angiogenesis Induced by Growth Factors

Angiogenesis is induced in the *in vivo* rabbit eye model assay with growth factors bFGF or VEGF and is described in the following sections.

a. Preparation of Hydron Pellets Containing Growth Factor and Monoclonal Antibodies

Hydron polymer pellets containing growth factor are prepared as described by D'Amato, et al., Proc. Natl. Acad. Sci., USA, 91:4082-

4085 (1994). The individual pellets contain 650 ng of the growth factors separately bound to sucralfate (Carafet, Marion Merrell Dow Corporation) to stabilize the growth factor and ensure its slow release into the surrounding tissue. In addition, hydron pellets are prepared containing a desired Src-expressing

5 retrovirus as previously described. The pellets are cast in specially prepared Teflon pegs that have a 2.5 mm core drilled into their surfaces. Approximately 12 ul of casting material is placed into each peg and polymerized overnight in a sterile hood. Pellets are then sterilized by ultraviolet irradiation. Effects of Src proteins are then assessed as previously described.

10 6. In Vivo Regression of Tumor Tissue Growth With Src Modulators As Measured by Chimeric Mouse:Human Assay

An in vivo chimeric mouse:human model is generated by replacing a portion of skin from a SCID mouse with human neonatal foreskin. The in vivo chimeric mouse:human model is prepared essentially as described in Yan, et al., J. Clin. Invest., 91:986-996 (1993). Briefly, a 2 cm² square area of skin is surgically

15 removed from a SCID mouse (6-8 weeks of age) and replaced with a human foreskin. The mouse is anesthetized and the hair removed from a 5 cm² area on each side of the lateral abdominal region by shaving. Two circular graft beds of 2 cm² are prepared by removing the full thickness of skin down to the fascia. Full

20 thickness human skin grafts of the same size derived from human neonatal foreskin are placed onto the wound beds and sutured into place. The graft is covered with a Band-Aid which is sutured to the skin. Micropore cloth tape is also applied to cover the wound.

The M21-L human melanoma cell line or MDA 23.1 breast carcinoma cell

25 line (ATCC HTB 26; $\alpha_v\beta_3$ negative by immunoreactivity of tissue sections with mAb LM609), are used to form the solid human tumors on the human skin grafts on the SCID mice. A single cell suspension of 5×10^6 M21-L or MDA 23.1 cells is injected intradermally into the human skin graft. The mice are then observed for 2 to 4 weeks to allow growth of measurable human tumors.

30 After a measurable tumor is established, retrovirus preparations of the present invention or PBS is injected into the mouse tail vein. Following a 2-3 week period, the tumor is excised and analyzed by weight and histology. The effect of expressed Src proteins of the present invention on the tumors is then assessed.

7. In Vitro Regression of Human Tumor Tissue Growth With Src Modulators As Measured by CAM Assay

5 Tumor growth depends on angiogenesis (Folkman, 1992; Weidner et al., 1991; Brooks et al., 1994b). In fact, recent reports suggest that tumor growth is susceptible to the anti-angiogenic effects of VEGF receptor antagonists (Kim et al., 1993). Therefore, we examined whether suppression of angiogenesis by delivery of kinase-deleted Src 251 would influence the growth of a human medulloblastoma (DAOY), a highly angiogenic tumor known to produce VEGF and very little bFGF (data not shown).

10 The 3 and 6 day DAOY medulloblastoma tumor growth assays were performed in the chick CAM essentially as previously described (Brooks et al., 1994). 5×10^6 DAOY cells cultured in RPMI 1640 containing 10% fetal calf serum were washed and seeded on the CAM of a 10 day embryo to produce DAOY tumor fragments. After 7 days 50 mg tumor fragments were dissected and
15 reseeded on another 10 day embryo and incubated for another 3 or 6 days with the topical application (25 μ l) of either control RCAS-GFP retrovirus, RCAS-Src 251, or mock treatment. Using the whole tissue confocal imaging of infected tumors as a guide we were able to determine that there was significant expression of the RCAS constructs around and within the tumor fragment with this topical approach.
20 Tumor resections and weighing were performed in a double blind manner removing only the easily definable solid tumor mass (Brooks et al., 1994). The wet tumor weights after 3 or 6 days were compared with initial weight and the percent change of tumor weight determined for each group.

25 These tumors readily grow on the CAM and produces active angiogenesis (FIG. 9) allowing us to selectively target the avian-derived tumor vasculature by using an avian-specific RCAS retrovirus.

FIG. 9 depicts results that show retroviral delivery of RCAS-Src 251 to human tumors growing on the chick CAM reverses tumor growth. FIG. 9A shows human medulloblastomas that were grown on the CAM of chick embryos as described above. Retrovirus containing RCAS-GFP or RCAS-Src 251 was
30 topically applied to preestablished tumors of greater than 50 mg. A representative micrograph of a medulloblastoma tumor fragment infected with RCAS-GFP expressing GFP reveals exclusive expression in the tumor blood vessels

(arrowhead) as detected by optical sectioning with a Bio Rad laser confocal scanning microscope (bar=500 μ m). FIG. 9B shows results from tumors treated as above that were allowed to grow for 3 or 6 days after which they were resected and wet weights determined. Data are expressed as the mean change in tumor weight (from the 50 mg tumor starting weight) +/- SEM of 2 replicates. RCAS-Src 251 had a significant impact on tumor growth after 3 days (*, $P < 0.002$) and 6 days (**, $P < 0.05$). FIG. 9C shows representative stereomicrographs of medulloblastoma tumors surgically removed from the embryos were taken with an Olympus stereomicroscope (bar=350 μ m). (Lower panel) A high magnification micrograph of each tumor showing the vasculature of each tumor in detail (bar=350 μ m). The arrowhead indicates blood vessel disruption in RCAS-Src251-treated tumors.

The results show that delivery of RCAS containing Src 251 to preestablished medulloblastomas resulted in selective viral expression in the tumor-associated blood vessels (FIG. 9A) and this ultimately led to the regression of these tumors within the span of six days (FIG. 9B). Importantly, the tumor-associated blood vessels in animals treated with virus containing Src 251 were severely disrupted and fewer in number compared to the tumor vessels in control animals (FIG. 9C). The fact that RCAS-GFP infected tumors showed GFP localization only in the tumor vasculature suggests that the anti-tumor effects observed with retrovirally delivered Src 251 were due to its anti-angiogenic properties.

The foregoing examples and the accompanying description are illustrative, and are not to be taken as limiting. The present invention also is not to be limited in scope by the cell line deposited, since the deposited embodiment is intended as a single illustration of one aspect of the invention. Any cell line that is functionally equivalent is within the scope of this invention. The deposit of material does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustration that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

What Is Claimed Is:

1. An article of manufacture comprising packaging material and a pharmaceutical composition contained within said packaging material, wherein said pharmaceutical composition is capable of modulating angiogenesis in a tissue associated with a disease condition, wherein said packaging material comprises a label which indicates that said pharmaceutical composition can be used for treating disease conditions by modulating angiogenesis, and wherein said pharmaceutical composition comprises a Src protein or an oligonucleotide having a nucleotide sequence capable of expressing said protein.
2. The article of manufacture of claim 1 wherein said Src protein is an active Src protein and said modulating potentiates angiogenesis.
3. The article of manufacture of claim 2 wherein said active Src protein is Src A.
4. The article of manufacture of claim 2 wherein said tissue has poor circulation.
5. The article of manufacture of claim 1 wherein said tyrosine kinase Src protein is an inactive Src protein and said modulating inhibits angiogenesis.
6. The article of manufacture of claim 5 wherein said inactive Src protein is Src 251 or Src K295M.
7. The article of manufacture of claim 5 wherein said tissue is inflamed and said condition is arthritis or rheumatoid arthritis.
8. The article of manufacture of claim 5 wherein said tissue is a solid tumor or solid tumor metastasis.
9. The article of manufacture of claim 8 wherein said administering is conducted in conjunction with chemotherapy.
10. The article of manufacture of claim 5 wherein said tissue is retinal tissue and said condition is retinopathy, diabetic retinopathy or macular degeneration.
11. The article of manufacture of claim 5 wherein said tissue is at the site of coronary angioplasty and said condition is restenosis.
12. The article of manufacture of claim 1 wherein said administering comprises intravenous, transdermal, intrasynovial, intramuscular, or oral administration.

13. The article of manufacture of claim 1 wherein said administering comprises a single dose intravenously.

14. The article of manufacture of claim 1 wherein said pharmaceutical composition further comprises a liposome.

5 15. The article of manufacture of claim 1 wherein said pharmaceutical composition comprises a viral expression vector capable of expressing said nucleotide sequence.

16. The article of manufacture of claim 1 wherein said pharmaceutical composition comprises an non-viral expression vector capable of expressing said
10 nucleotide sequence.

17. A method for modulating angiogenesis in a tissue associated with a disease condition comprising administering to said tissue a pharmaceutical composition comprising a Src protein or a nucleotide sequence capable of expressing said protein.

15 18. The method of claim 17 wherein said Src protein is an active Src protein and said modulating potentiates angiogenesis.

19. The method of claim 18 wherein said active Src protein is Src A.

20. The method of claim 18 wherein said tissue has poor circulation.

21. The method of claim 17 wherein said Src protein is an inactive Src
20 protein and said modulating inhibits angiogenesis.

22. The method of claim 21 wherein said inactive Src protein is Src 251 or Src K295M.

23. The method of claim 21 wherein said tissue is inflamed and said condition is arthritis or rheumatoid arthritis.

25 24. The method of claim 21 wherein said tissue is a solid tumor or solid tumor metastasis.

25. The method of claim 24 wherein said administering is conducted in conjunction with chemotherapy.

26. The method of claim 21 wherein said tissue is retinal tissue and said
30 condition is retinopathy, diabetic retinopathy or macular degeneration.

27. The method of claim 21 wherein said tissue is at the site of coronary angioplasty and said tissue is at risk for restenosis.

28. The method of claim 17 wherein said administering comprises intravenous, transdermal, intrasynovial, intramuscular, or oral administration.

29. The method of claim 17 wherein said administering comprises a single dose intravenously.

5 30. The method of claim 17 wherein said pharmaceutical composition further comprises a liposome.

31. The method of claim 17 wherein said pharmaceutical composition comprises an retroviral expression vector capable of expressing said nucleotide sequence.

10 32. The method of claim 17 wherein said pharmaceutical composition comprises an non-viral expression vector capable of expressing said nucleotide sequence.

33. A pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a viral gene transfer vector containing a nucleic acid
15 and pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein, said src protein having any amino acid residue at codon 527 except for tyrosine, serine or threonine.

34. A pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a non-viral gene transfer vector containing a nucleic
20 acid and pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein and said src protein having any amino acid residue at codon 527 except tyrosine, serine or threonine.

35. A pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a viral gene transfer vector containing a nucleic acid
25 and pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein having no kinase activity.

36. A pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a non-viral gene transfer vector containing a nucleic
30 acid and pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein, said src protein having no kinase activity.

37. A pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a therapeutic amount of a src protein in a

pharmaceutically acceptable carrier or excipient; said src protein having any amino acid residue at codon 527 except tyrosine, serine or threonine.

38. A pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a a src protein in a pharmaceutically acceptable carrier or excipient; said src protein having no kinase activity.
- 5

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

09701500 112900



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|--|------------------|---|
| <p>(51) International Patent Classification ⁶ : C12N 9/00</p> | <p>A1</p> | <p>(11) International Publication Number: WO 99/61590 (43) International Publication Date: 2 December 1999 (02.12.99)</p> |
| <p>(21) International Application Number: PCT/US99/11780 (22) International Filing Date: 28 May 1999 (28.05.99) (30) Priority Data: 60/087,220 29 May 1998 (29.05.98) US (71) Applicant (for all designated States except US): THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10550 North Torrey Pines Road, La Jolla, CA 92037 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CHERESH, David, A. [US/US]; 327 Via Andalusia, Encinitas, CA 92024 (US). ELICEIRI, Brian [US/US]; 935 Laguna #5, Carlsbad, CA 92008 (US). SCHWARTZBERG, Pamela, L. [US/US]; 49/4A38, National Humane Genome Research Institute, National Institute of Health, Bethesda, MD 20892-4472 (US). (74) Agents: CEPURITIS, Talivaldis et al.; Olson & Hierl, Ltd., 36th floor, 20 North Wacker Drive, Chicago, IL 60606 (US).</p> | | <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> |
| <p>(54) Title: METHODS AND COMPOSITIONS USEFUL FOR MODULATION OF ANGIOGENESIS USING TYROSINE KINASE SRC</p> | | |
| <p>(57) Abstract The present invention describes methods for modulating angiogenesis in tissues using Src protein, modified Src protein, and nucleic acids encoding for such. Particularly the invention describes methods for inhibiting angiogenesis using an inactive Src protein, or nucleic acids encoding therefor, or for potentiating angiogenesis using an active Src protein, or nucleic acids encoding therefor. The invention also describes the use of gene delivery systems for providing nucleic acids encoding for the Src protein, or modified forms thereof.</p> | | |

CHICKEN c-SRC cDNA

(SEQ ID NO:2)

1 tctgacaccc atctgtctgt ctgtctgtgt gctgcaggag ctgagctgac tctgtctgtg
61 cctcgcgtac cactgtggcc aggcggtagc tgggacgtgc agcccaccac catggggagc
121 agcaagagca agcccaagga ccccgaccag cgccggcgca gcctggagcc acccgacagc
181 acccaccacg ggggattccc agcctcgcag accccaaca agacagcagc ccccgacacg
241 caccgcaccc ccagccgctc ctttgggacc gtggccaccg agcccaagct ctccgggggc
301 ttcaaacatt ctgacaccgt tacgtgccg cagcgtgccg gggcactggc tggcggcgctc
361 accactttcg tggctctcta cgactacgag tcccggactg aaacggactt gtcctcaag
421 aaaggagaac gctgcagat tgtaacaac acggaagggtg actggtggct ggctcattcc
481 ctactacag gacagacggg ctacatcccc agtaactatg tcgcgccctc agactccatc
541 caggctgaag agtggtaact tgggaagatc actcgtcggg agtccgagcg gctgtctctc
601 aaccccgaaa acccccgggg aacctcttg gtccgggaga gcgagacgac aaaagggtgcc
661 tattgectct ccgtttctga ctttgacaac gccaaagggc tcaatgtgaa gcactacaag
721 atccgcaagc tggacagcgg cggtctctac atcacctcac gcacacagtt cagcagcctg
781 cagcagctgg tggcctacta ctccaaacat gctgatggct tgtgccaccg cctgaccaac
841 gtctgccccg cgtccaagcc ccagaccag ggactcgcca aggacgcgtg ggaaatcccc
901 cgggagtcgc tgcggctgga ggtgaagctg gggcagggct gctttggaga ggtctggatg
961 gggacctgga acggcaccac cagagtggcc ataaagactc tgaagcccg gaccatgtcc
1021 ccggaggcct tcctgcagga agcccaagtg atgaagaagc tccggcatga gaagctgggt
1081 cagctgtacg cagtgggtgc ggaagagccc atctacatcg tctctgagta catgagcaag
1141 gggagcctcc tggatttctt gaaggagag atgggcaagt acctgcggct gccacagctc
1201 gtcgatatgg ctgctcagat tgcattccgg atggcctatg tggagaggat gaactacgtg
1261 caccgagacc tgcgggcggc caacatcctg gtgggggaga acctggtgtg caaggtggct
1321 gactttgggc tggcacgcct catcaggagc aacaggtaca cagcacggca aggtgccaag
1381 ttcccatca agtggacagc ccccgaggca gccctctatg gccgggtcac catcaagtcg
1441 gatgtctggt ccttcggcat cctgtgact gagctgacca ccaagggccg ggtgccatac
1501 ccagggatgg tcaacaggga ggtgctggac caggtggaga ggggctaccg catgccctgc
1561 ccgcccagat gcccagatc gtgcatgac ctcatgtgcc agtgcctggc gagggaccct
1621 gaggagcggc ccactttga gtacctgag gccttctgg aggactacti cacctcgaca
1681 gagccccagt accagcctgg agagaaccta taggcctgga gtcctctcg gaccagaggc
1741 ctgcgtgtgg ggtacaggg

FIG. 1

CHICKEN cSRC ENCODED PROTEIN

(SEQ ID NO:3)

MGSSKSKPKDPSQRRRSLEPPDSTHHGGFPASQTPNKTA
PDTHRTSPRSFGTVATEPKLFGGFNTSDTVTSPQRAGALA
GGVTTFVALYDYESTETDLSFKKGERLQIVNNTSGDWL
AHSLTTGQTGYIPSNYVAPSDSIQAEWYFGKITRRESER
LLNPENPRGTFLVRESETTKGA YCLSVSDFDNAKGLNVK
HYKIRKLDSGGFYITSRTQFSSLQQLVAYYSKHADGLCHR
LTNVCPTSKPQTQGLAKDAWEIPRESLRLEVKLGGGCFGE
VWMGTWNGTTRVAIKTLKPGTMSPEAFLQEAQVMKKLRHE
KLVQLYAVVSEEPYIVTEYMSKGSLLDFLKGEMGKYLRL
PQLVDMAAQIASGMAYVERMNYVHRDLRAANILVGENL
VCKVADFGGLARLIEDNEYTARQGAKFPIKWTAPAAALYGR
FTIKSDVWSFGILLTELTTKGRVPYPGMVNREVLDQVERG
YRMPCPPECPESLHDLMCQCWRRDPEERPTFEYLQAFLE
DYFTSTEPQYQPGENL

FIG. 2

3/10

HUMAN c-SRC cDNA

(SEQ ID NO:4)

1 gcgccgcgtc ccgcaggccg tgatgccgc cgcgcggagg tggcccggac cgcagtgcc
61 caagagagct ctaatggtac caagtacag gttggcttta ctgtgactcg gggacgccag
121 agctcctgag aagatgtcag caatacaggc cgctggcca tccggtacag aatgtattgc
181 caagtacaac ttccacggca ctgccgagca ggacctgccc ttctgcaaag gagacgtgct
241 caccattgtg gccgtcacca aggaccccaa ctggtacaaa gccaaaaaca aggtgggccc
301 tgagggcatc atcccagcca actacgtcca gaagcgggag ggctgaagg cgggtacca
361 actcagcctc atgccttggt tccacggcaa gatcacacgg gagcaggctg agcggcttct
421 gtacccgccg gagacaggcc tgttctggt gcgggagagc accaactacc ccggagacta
481 cacgctgtgc gtgagctgcg acggcaagggt ggagcactac cgcatcatgt accatgccag
541 caagctcagc atcgacgagg aggtgtactt tgagaacctc atgcagctgg tggagcacta
601 cacctcagac gcagatggac tctgtacgc cctcattaaa ccaaagggtca tggagggcac
661 agtggcggcc caggatgagt tctaccgcag cggtgggccc ctgaacatga aggagctgaa
721 gctgctgcag accatcgga agggggaggt cggagacgtg atgctggcg attaccgagg
781 gaacaaagtc gccgtcaagt gcattaagaa cgacgccact gccaggcct tcttggtga
841 agcctcagtc atgacgcaac tgcggcatag caacctggtg cagctcctgg gcgtgatcgt
901 ggaggagaag ggcgggctct acatcgtcac tgagtacatg gccaggggga gccttgtga
961 ctacctgcgg tctaggggtc ggtcagtgct gggcggagac tgtctctca agttctcgt
1021 agatgtctgc gaggccatgg aatacctgga gggcaacaat ttcgtgcatc gagacctggc
1081 tgcccgaat gtgtggtgt ctgaggacaa cgtggccaag gtcagcgact ttggtctac
1141 caaggaggcg tccagcacc aggacacggg caagctgcca gtcaagtga cagcccctga
1201 gggcctgaga gagaagaaat tctcactaa gtctgacgtg tggagtctc gaatcctct
1261 ctgggaaatc tactccttg ggcgagtgcc ttatcaaga attcccctga aggacgtcgt
1321 ccctcgggtg gagaagggt acaagatgga tgccccgcac ggctgcccgc ccgcagtcta
1381 tgaagtcatg aagaactgct ggcacctgga cgcgcctatg cggccctcct tctacagct
1441 ccgagagcag cttgagcaca tcaaaccca cgagctgcac ctgtgacggc tggcctccgc
1501 ctgggtcatg ggcctgtggg gactgaacct ggaagatcat ggacctggtg ccctgctca
1561 ctgggcccga gcctgaactg agccccagcg ggctggcggg ccttttct gcgtcccagc
1621 ctgcaccct ccggccccgt ctctcttga ccacctgtg gggcctgggg agcccactga
1681 ggggccaggg aggaaggagg ccacggagcg ggaggcagcg cccaccacg tcgggcttcc
1741 ctggcctccc gccactcgcc ttcttagagt ttattcctt tctttttg agattttt
1801 tccgtgtgtt ttttttat tttttcaa gataaggaga aagaaagta cagcaaat
1861 ggcattttac aagaagtacg aatctattt ttctgtctt gccgtgagg gtggggggga
1921 ccgggccccct cttagggac cctcgcccc agcctcatt cccattctgt gtcccatgtc
1981 ccgtgtctcc tcggtcgccc cgtgttgcc ctgacctg ttgactgtt tgcatcgcc
2041 cgaggcagac gtctgtcagg ggcttgatt tcgtgtgcc ctgccaccg cccaccgc
2101 ttgtgagatg gaattgtaat aaaccacgcc atgaggacac gcggcccgcc ctggcgctt
2161 cctccaccga aaaaaaaaa aaaaaaa

FIG. 3

4 / 10


HUMAN c-SRC ENCODED PROTEIN

(SEQ ID NO:5)



MSAIQAAWPSGTECIAKYNFHGTAEQDLPFCKGDVLTIVAVTKD
PNWYKAKNKVGREGIIPANYVQKREGVKAGTKLSLMPWFHGKIT
REQAERLLYPPETGLFLVRESTNYPGDYTLCVSCDGKVEHYRIMY
HASKLSIDEEVYFENLMQLVEHYTSDADGLCTRLIKPKVMEGTVA
AQDEFYRSGWALNMKELKLLQTIGKGEFGDVMLGDYRGNKVAV
KCIKNATAQAFLAEASVMTQLRHSNLVQLLGVIVEEKGGLYIVTE
YMAKGSLVDYLRSRGRSVLGGDCLLKFSLDVCEAMEYLEGNFVH
RDLAARNVLVSEDNVAKVSDFGLTKEASSTQDTGKLPVKWTAPEAL
REKKFSTKSDVWSFGILLWEIYSFGRVPYPRIPLKD VVPRVEKGYKM
DAPDGCPPAVYEVMKNCWHLDAAMRPSFLQLREQLEHIKTHELHL

FIG. 4


Activation of endogenous Src activity by bFGF and VEGF

**FIG. 5**



Effect of RCAS-mediated expression of
Src A on angiogenesis in the chick CAM

FIG. 6A**FIG. 6B**

Retroviral expression of Src A activates vascular MAP kinase phosphorylation



Mock



Src A





FIG. 7

8/10

Selective requirement for Src
activity during VEGF, but not
bFGF-induced angiogenesis




FIG. 8A





FIG. 8B

MOCK Src 251-myc

BLOT:ANTI-Src



- Src 251

FIG. 8C

9/10




FIG. 9A





FIG. 9B



GFP

Src 251

FIG. 9C

10/10





FIG. 10

Please type a plus sign (+) inside this box → 

Approved for use through 9/30/00. OMB 0651-0032
 Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

| | | |
|--|--------------------------|-------------|
| DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION (37 CFR 1.63) <input type="checkbox"/> Declaration Submitted with Initial Filing OR <input type="checkbox"/> Declaration Submitted after Initial Filing (surcharge (37 CFR 1.16 (e)) required) | Attorney Docket Number | TSRI 651.1 |
| | First Named Inventor | Cheresh |
| | COMPLETE IF KNOWN | |
| | Application Number | / |
| | Filing Date | 28 MAY 1999 |
| | Group Art Unit | |
| | Examiner Name | |

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Methods and Compositions Useful for Modulation of Angiogenesis
Using Tyrosine Kinase Src

the specification of which

(Title of the Invention)

☐ is attached hereto
OR

☒ was filed on (MM/DD/YYYY) 05/28/1999 as United States Application Number or PCT International

Application Number PCT/US99/11780 and was amended on (MM/DD/YYYY) (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

| Prior Foreign Application Number(s) | Country | Foreign Filing Date (MM/DD/YYYY) | Priority Not Claimed | Certified Copy Attached? | |
|-------------------------------------|---------|----------------------------------|--------------------------|--------------------------|--------------------------|
| | | | | YES | NO |
| | | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| | | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| | | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| | | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.

| Application Number(s) | Filing Date (MM/DD/YYYY) | |
|-----------------------|--------------------------|--|
| 60/087,220 | 05/29/1998 | <input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto. |

(Page 1 of 2)

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box → ☐ +

PTO/SB/01 (12-97)
 Approved for use through 9/30/00. OMB 0651-0032
 Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION — Utility or Design Patent Application

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

| U.S. Parent Application or PCT Parent Number | Parent Filing Date (MM/DD/YYYY) | Parent Patent Number (if applicable) |
|--|---------------------------------|--------------------------------------|
| | | |

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

☐ Customer Number

OR

☐ Registered practitioner(s) name/registration number listed below

Place Customer Number Bar Code Label here

| Name | Registration Number | Name | Registration Number |
|-------------------|---------------------|-----------------------|---------------------|
| Arne M. Olson | 30,203 | Michael A. Hierl | 29,807 |
| Dolores T. Kenney | 31,269 | Talivaldis Cepuritis | 20,818 |
| Seymour Rothstein | 19,369 | Daniel J. Deneufbourg | 33,675 |

☒ Additional registered practitioner(s) named on supplemental Registered Practitioner Information sheet PTO/SB/02C attached hereto.


Direct all correspondence to: ☐ Customer Number or Bar Code Label OR ☒ Correspondence address below

| | | | |
|---------|--|-----------|-----------------------|
| Name | <u>TALIVALDIS CEPURITIS</u> | | |
| Address | <u>OLSON & HIERL, LTD.</u> | | |
| Address | <u>20 NORTH WACKER DRIVE, 36TH FLOOR</u> | | |
| City | <u>CHICAGO</u> | State | <u>IL</u> |
| Country | <u>US</u> | Telephone | <u>(312) 580-1180</u> |
| | | Fax | <u>(312) 580-1189</u> |

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

| | | | |
|--|--------------------------|---|----------------|
| Name of Sole or First Inventor: | | <input type="checkbox"/> A petition has been filed for this unsigned inventor | |
| Given Name (first and middle (if any)) | | Family Name or Surname | |
| <u>David A.</u> | | <u>Cheresh</u> | |
| Inventor's Signature | <u>David A. Cheresh</u> | Date | <u>6/22/99</u> |
| Residence: City | <u>Encinitas</u> | State | <u>CA</u> |
| | | Country | <u>US</u> |
| Post Office Address | <u>327 Via Andalusia</u> | | |
| Post Office Address | | | |
| City | <u>Encinitas</u> | State | <u>CA</u> |
| | | ZIP | <u>92024</u> |
| | | Country | <u>US</u> |

☒ Additional inventors are being named on the 1 supplemental Additional Inventor(s) sheet(s) PTO/SB/02A attached hereto

Please type a plus sign (+) inside this box → PTO/SB/02A (3-97)
Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION**ADDITIONAL INVENTOR(S)**
Supplemental Sheet
Page 1 of 1

| | | | | | | | |
|---|--------------------------------|-------|-----------|---|--------------|-------------|----------------|
| Name of Additional Joint Inventor, if any: | | | | <input type="checkbox"/> A petition has been filed for this unsigned inventor | | | |
| Given Name (first and middle [if any]) | | | | Family Name or Surname | | | |
| <u>2-00</u> <u>Brian</u> | | | | <u>Eliceiri</u> | | | |
| Inventor's Signature | <u>Brian</u> | | | <u>elm</u> | | Date | <u>6/22/99</u> |
| Residence: City | <u>Carlsbad</u> | State | <u>CA</u> | Country | <u>US</u> | Citizenship | <u>US</u> |
| Post Office Address | <u>3104 Hataca Road</u> | | | | | | |
| Post Office Address | | | | | | | |
| City | <u>Carlsbad</u> | State | <u>CA</u> | ZIP | <u>92009</u> | Country | <u>US</u> |
| Name of Additional Joint Inventor, if any: | | | | <input type="checkbox"/> A petition has been filed for this unsigned inventor | | | |
| Given Name (first and middle [if any]) | | | | Family Name or Surname | | | |
| <u>3-00</u> <u>Pamela L.</u> | | | | <u>Schwartzberg</u> | | | |
| Inventor's Signature | <u>Pamela L. Schwartzberg</u> | | | <u>9-6-99</u> | | Date | <u>9-6-99</u> |
| Residence: City | <u>Bethesda</u> | State | <u>MD</u> | Country | <u>US</u> | Citizenship | <u>US</u> |
| Post Office Address | <u>5521 Spruce Tree Avenue</u> | | | | | | |
| Post Office Address | | | | | | | |
| City | <u>Bethesda</u> | State | <u>MD</u> | ZIP | <u>20814</u> | Country | <u>US</u> |
| Name of Additional Joint Inventor, if any: | | | | <input type="checkbox"/> A petition has been filed for this unsigned inventor | | | |
| Given Name (first and middle [if any]) | | | | Family Name or Surname | | | |
| | | | | | | | |
| Inventor's Signature | | | | | | Date | |
| Residence: City | | State | | Country | | Citizenship | |
| Post Office Address | | | | | | | |
| Post Office Address | | | | | | | |
| City | | State | | ZIP | | Country | |

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box → ☐

PTO/SB/02C (3-97)

Approved for use through 9/30/98. OMB 0651-0032

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

| DECLARATION | | REGISTERED PRACTITIONER INFORMATION (Supplemental Sheet) | |
|--------------------|---------------------|---|---------------------|
| Name | Registration Number | Name | Registration Number |
| Steven D. Weseman | 41,372 | | |
| Timothy L. Harney | 38,174 | | |
| Mark Chao | 37,293 | | |
| Joseph M. Kuo | 38,943 | | |
| James C. Haight | 25,588 | | |
| David R. Sadowski | 32,808 | | |
| Robert Benson | 33,612 | | |
| Jack Spiegel | 34,477 | | |
| Susan S. Rucker | 35,762 | | |
| Steven M. Ferguson | 38,448 | | |
| Stephen L. Finley | 36,357 | | |
| John Peter Kim | 38,514 | | |

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

09/701500

1/20

SEQUENCE LISTING

<110> CHERESH, David A.
ELICEIRI, Brian
SCHWARTZBERG, Pamela L.

<120> METHODS AND COMPOSITIONS USEFUL FOR MODULATION OF
ANGIOGENESIS USING TYROSINE KINASE SRC

<130> TSRI 651.1

<140> US 09/701,500
<141> 2000-11-29

<150> PCT/US99/11780
<151> 1999-05-28

<150> US 60/087,220
<151> 1998-05-29

<160> 6

<170> PatentIn Ver. 2.0

<210> 1
<211> 11627
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:RCASBP(A) based
on avian sarcoma virus

<220>
<221> misc_feature
<222> (7649)..(11258)
<223> pBR322 sequences

<220>
<221> LTR
<222> (7166)..(7494)
<223> upstream

<220>
<221> LTR
<222> (1)..(101)
<223> upstream (numbering begins at the upstream R)

<220>
<221> misc_feature
<222> (11394)..(11623)
<223> U3

<220>
<221> misc_feature
<222> (1)..(21)

2/20

<223> R

<220>

<221> misc_feature

<222> (22)..(101)

<223> U5

<220>

<221> misc_feature

<222> (102)..(119)

<220>

<221> LTR

<222> (7166)..(7494)

<223> downstream

<220>

<221> misc_feature

<222> (7166)..(7393)

<223> U3

<220>

<221> misc_feature

<222> (7394)..(7414)

<223> R

<220>

<221> misc_feature

<222> (7415)..(7494)

<223> U5

<220>

<221> misc_feature

<222> (7154)..(7165)

<223> PPT

<220>

<221> misc_feature

<222> (388)..(391)

<223> splice donor (AGGT)

<220>

<221> misc_feature

<222> (5074)..(5077)

<223> env splice acceptor (AGGC)

<220>

<221> misc_feature

<222> (6982)..(6985)

<223> ClaI splice acceptor (AGGA)

3/20

<220>
<221> gene
<222> (372)..(902)
<223> gag p19

<220>
<221> gene
<222> (909)..(1094)
<223> gag p10

<220>
<221> gene
<222> (1095)..(1814)
<223> gag p27

<220>
<221> gene
<222> (1843)..(2108)
<223> gag p12

<220>
<221> gene
<222> (2109)..(2480)
<223> gag p15

<220>
<221> misc_signal
<222> (2481)..(2483)
<223> gag stop

<220>
<221> gene
<222> (2501)..(4216)
<223> pol RT

<220>
<221> gene
<222> (4217)..(5185)
<223> pol IN

<220>
<221> misc_signal
<222> (5186)..(5188)
<223> pol stop

<220>
<221> gene
<222> (5244)..(6263)
<223> env gp85

4/20

<220>
<221> gene
<222> (6264)..(6878)
<223> env gp37

<220>
<221> misc_signal
<222> (6879)..(6881)
<223> env stop

<220>
<221> misc_feature
<222> (7027)
<223> ClaI site/ the ClaI site in gag is methylated in
Dam+ strains and does not cut.

<400> 1
gccatttgac cattcaccac attggtgtgc acctgggttg atggccggac cgttgattcc 60
ctgacgacta cgagcacctg catgaagcag aaggcttcat ttggtgaccc cgacgtgata 120
gttagggaat agtggtcggc cacagacggc gtggcgatcc tgtctccatc cgtctcgtct 180
atcgggaggc gagttcgatg accctggttg agggggctgc ggcttaggga ggcagaagct 240
gagtaccgtc ggagggagct ccagggcccc gagcgactga cccctgccga gaactcagag 300
ggtcgtcgga agacggagag tgagcccgac gaccaccca ggcacgtctt tggtcggcct 360
gcggatcaag catggaagcc gtcattaagg tgatttcgtc cgcggtgtaa acctattgct 420
ggaaaatctc tccttctaag aaggaaatcg gggccatgtt gtccctgtta caaaaggaag 480
ggttgcttat gtctccctca gatttatatt ctccggggtc ctgggatccc atcactgcgg 540
cgctctccca gcgggcaatg gtacttgga aatcgggaga gttaaaaacc tggggattgg 600
ttttgggggc attgaaggcg gctcgagagg aacagggttac atctgagcaa gcaaagtttt 660
ggttgggatt agggggaggg agggctctct ccccaggctc ggagtgcac gagaaaccag 720
ctacggagcg gcgaatcgac aaaggggagg aggtgggaga aacaactgtg cagcgagatg 780
cgaagatggc gccagaggaa gcgccacac ctaaaaccgt tggcacatcc tgctatcatt 840
gcggaacagc tgttggtgc aattgcgcca ccgccacagc ctcgcccct cctccccctt 900
atgtggggag tggtttgtat ccttccctgg cgggggtggg agagcagcag ggccagggag 960
ataacacgtc tcggggggcg gacgagccaa gggaggagcc agggcacgcg ggtcaggccc 1020
ctgggccggc cctgactgac tgggcaagg taaggaggga gcttgcgagt actggtccgc 1080
ccgtggtggc catgcctgta gtgattaaga cagagggacc cgctggacc cctctggagc 1140
caaaattgat cacaagactg gctgatacgg tcaggaccaa gggcttacga tccccgatca 1200

ctatggcaga agtgggaagcg ctcatgtcct ccccgttgct gcgcgatgac gtcacgaatc 1260
taatgagagt gatttttagga cctgccccat atgccttatg gatggacgct tggggagtc 1320
aactccagac ggttatagcg gcagccactc gcgacccccg acaccacagcg aacgggtcaag 1380
ggcgggggga acggactaac ttggatcgat taaagggcct agctgatggg atggtgggca 1440
accacaggg tcaggccgca ttattaagac cgggggaatt ggttgctatt acggcgctcg 1500
ctctccaggc gtttagagaa gttgcccggc tggcgggaacc tgcaggtcca tgggcggaca 1560
tcacgcaggg accatctgag tcctttgttg attttgccaa tggccttata aaggcggttg 1620
aggggtcaga tctccgcct tccgcgcggg ctccggtgat cattgactgc tttaggcaga 1680
agtcacagcc agatattcag cagcttatac gggcagcacc ctccacgctg accaccccag 1740
gagagataat caaatatgtg ctagacaggc agaagattgc ccctcttacg gatcaaggca 1800
tagccgcggc catgtcgtct gctatccagc ccttagttat ggcagtagtc aatagagaga 1860
gggatggaca aactgggtcg ggtggtcgtg cccgagggct ctgctacact tgtggatecc 1920
cgggacatta tcaggcacag tgcccgaaaa aacgaaagtc aggaaacagc cgtgagcgat 1980
gtcagctgtg tgacgggatg ggacacaacg ctaaacagt taggaagcgg gatggcaacc 2040
agggccaacg cccaggaaga ggtctctctt cggggccgtg gcccggccct gagcagcctg 2100
ccgtctcggt agcgatgaca atggaacata aagatcgccc cttggttagg gtcattctga 2160
ctaactctgg gagtcatcca gtcaaacaac gttcgggtga tatcaccgcg ctggttgact 2220
ccggagcgga catcactatt atttcggagg aggattggcc tactgattgg ccggtggtgg 2280
acaccgcgaa cccacagatc catggcatag gagggggaat tcccatgca aaatcccggg 2340
atatgataga ggtgggggtt attaaccgag acgggtcgtt ggagcgaccc ctgctcctct 2400
tccccgcagt cgctatggtt agaggagta tcctaggaag agattgtctg cagggcctag 2460
ggctccgctt gacaaattta tagggagggc cactgttctc actgttgccg tacatctggc 2520
tattccgctc aaatggaagc cagaccgcac gcctgtgtgg attgaccagt ggccccctcc 2580
tgaaggtaaa cttgtaggcc taacgcaatt agtggaaaaa gaattacagt taggacatat 2640
agagccctca cttagttggt ggaacacacc tgtttttcgt gatccggaag gcttccgggt 2700
cttatcgctt attgcatgat ttgcgcgctg ttaacgccaa gcttgctcct tttggggccg 2760
tccaacaggg ggcgccagtt ctctccgcgc tcccgcgtgg ctggccccctg atggtcctag 2820
acctcaagga ttgcttcttt tctatccctc ttgcggaaca agatcgcgaa gcttttgc 2880
ttacgctccc ctctgtgaat aaccaggccc ccgctogaag attccaatgg aaggtcttgc 2940

cccaagggat gacctgttct ccactatct gtcagttggt agtgggtcag gtgctcgagc 3000
ccttgcgact caagcaccca gctctgcga tgttgatta tatggacgat cttttgctag 3060
ccgctcaag tcatgatggg ttggaagcgg caggaagga gggtatcggg acattggaaa 3120
gagccgggtt cactatttcg ccgataaga tccagagga gcccgagta caatatcttg 3180
ggtacaagtt aggcagtacg tatgtagcac ccgtaggctt ggtagcagaa ccaggatag 3240
ccacctgtg ggatgttcaa aagctggtg ggtcacttca gtggcttcgc ccagcgtag 3300
ggatcccgcc acgactgatg ggtccctttt atgagcagtt acgagggtca gatcctaacg 3360
aggcgagga atggaatcta gacatgaaaa tggcctggag agagatcgta cagcttagca 3420
ctactgctgc cttggaacga tgggaccctg ccagcctct ggaaggagcg gtcgctagat 3480
gtgaacaggg ggcaataggg gtctgggac agggactgtc cacacacca aggccatgtt 3540
tgtggttatt ctccaccaa ccaccaagg cgtttactgc ttggttagaa gtgctcacc 3600
ttttgattac taagctacgc gcttcggcag tgcgaacctt tggcaaggag gttgatatcc 3660
tcctgttgcc tgcattcttc cgggaggacc ttccgctccc ggaggggatc ctgtagcac 3720
ttagggggtt tgcaggaaaa atcaggagta gtgacacgcc atctatttt gacattgccc 3780
gtccactgca tgtttctctg aaagtgagg ttaccgacca ccctgtgccg ggacccactg 3840
tctttaccga cgctctctca agcaccata aaggggtggt agtctggagg gagggcccaa 3900
ggtgggagat aaaagaaata gttgatttgg gggcaagtgt acaacaactg gaggcacgcg 3960
ctgtggccat ggcattctg ctgtggcca caacgccac taatgtagt actgactctg 4020
cgtttggttc gaaaatgtta ctcaagatgg gacaggagg agtcccgct acagcgcg 4080
cttttatttt agaggatgag ttaagccaaa ggtcagccat ggccgccgtt ctccacgtgc 4140
ggagtcattc tgaagtcca gggtttttca cagaaggaaa tgacgtggca gatagccaag 4200
ccacctttca agcgtatccc ttgagagagg ctaaagatct tcataccgct ctccatattg 4260
gacccgcgc gctatccaaa gcgtgtaata tatctatgca gcaggctagg gaggttggtc 4320
agacctgcc gcattgtaat tcagccctg cgttgaggc cggggtaaac ctaggggtt 4380
tgggaccct acagatatgg cagacagact ttacgcttga gcctagaatg gctccccgtt 4440
cctggctgc tgttactgtg gacaccgct catcagcag agtcgtaact cagcatggc 4500
gtgttacatc ggttgctgca caacatcatt gggccacggc tategccgtt ttgggaagac 4560
caaaggccat aaaaacagat aacgggtcct gcttcacgtc cagatccacg cgagagtggc 4620
tcgcgagatg ggggtagca cacaccacg ggattccggg aaattcccag ggtcaagcta 4680

7/20

tggtagagcg ggccaaccgg ctctgaaag ataagatccg tgtgctcgcg gagggggacg 4740
gctttatgaa aagaatcccc accagcaaac aggggggaact attagccaag gcaatgtatg 4800
ccctcaatca ctttgagcgt ggtgaaaaca caaaaacacc gatacaaaaa cactggagac 4860
ctaccgttct tacagaagga ccccggtta aaatacgaat agagacaggg gagtgggaaa 4920
aaggatggaa cgtgctggtc tggggacgag gttatgccgc tgtgaaaaac agggacactg 4980
ataaggttat ttgggtaccc tctcgaaaag ttaaaccgga tgtcacccaa aaggatgagg 5040
tgactaagaa agatgaggcg agccctcttt ttgcaggcat ttctgactgg ataccctggg 5100
aagacgagca agaaggactc caaggagaaa ccgctagcaa caagcaagaa agacccgag 5160
aagacacctt tgctgccaac gagagttaat tatattctca ttattggtgt cctgggtcttg 5220
tgtgaggtta cgggggtaag agctgatgtc cacttactcg agcagccagg gaacctttgg 5280
attacatggg ccaaccgtac aggccaaacg gatttttggc tctctacaca gtcagccacc 5340
tccccctttc aaacatgttt gataggtatc ccgtccccta tttccgaggg tgattttaag 5400
ggatatgttt ctgatacaaa ttgcaccacc ttgggaactg atcggttagt ctcgtcagcc 5460
gactttactg ggggacctga caacagtacc accctcactt atcggaaggt ctcatgcttg 5520
ttgttaaagc tgaatgtctc tatgtgggat gagccacctg aactacagct gttaggttcc 5580
cagtctctcc ctaacattac taatattgct cagatttccg gtataaccgg gggatgcgta 5640
ggcttcagac cacaaggggt tccttggtat ctaggttggg ctagacagga ggccacgcg 5700
tttctcetta gacacccctc tttctctaaa tccacggaac cgtttacagt ggtgacagcg 5760
gataggcaca atctttttat ggggagttag tactgcggtg catatggcta cagattttgg 5820
aacatgtata actgctcaca ggtggggcg cagtaccgct gtggtaatgc ggcacgccc 5880
cgcacgggtc ttcctgaaat ccagtgtaca aggagaggag gcaaatgggt taatcaatca 5940
caggaaatta atgagtcgga gccgttcagc tttacgggtga actgtacagc tagtagtttg 6000
ggtaatgcca gtgggtggtg cggaaaagca ggcacgattc tcccgggaaa gtgggtcgac 6060
agcacacaag gtagtttcac caaaccaaaa gcgtaccac ccgcaatttt cctcatttgt 6120
ggggatcgcg catggcaagg aattcccagt cgctcggtag ggggccctg ctatttaggc 6180
aagcttacca tgtagcacc taagcataca gatattctca aggtgcttgt caattcatcg 6240
cggacaggta taagacgtaa acgaagcacc tcacacctgg atgatacatg ctcatatgaa 6300
gtgcagcttt ggggtcctac agcaagaatc tttgcatcta tcctagcccc gggggtagca 6360
gctgcgcaag ccttaagaga aattgagaga ctagcctggt ggtccgttaa acaggctaac 6420

8/20

ttgacaacat cactcctcgg ggacttattg gatgatgtca cgagtattcg acacgcggtc 6480
ctgcagaacc gagcggctat tgacttcttg ctccatagctc acggccatgg ctgtgaggac 6540
gttgccggaa tgtgctgttt caatttgagt gatcagagtg agtctataca gaagaagttc 6600
cagctaataga aggaacatgt caataagatc ggcgtggata gcgacctaat tggaagttgg 6660
ctgcgaggac tattcggggg aataggagaa tggggcgttc atttgctgaa aggactgctt 6720
ttggggcttg tagttatttt gttgctagta gtgtgcctgc cttgcctttt gcaaagtgtta 6780
tgcggttaata ggagaaagat gattaataac tccatcagct accacacgga atataagaag 6840
ctgcaaaagg cctgtgggca gcctgaaagc agaatagtat aaggcagtac atgggtggtg 6900
gtatagcgct tgcgagtcca tcgagcaagg caggaaagac agctattggg aattgtgaaa 6960
tacgcttttg tctgtgtgct gcaggagctg agctgactct gctggtggcc tcgcgtacca 7020
ctgtggcatc gatgcgatgt acggggcaga tatacgcgta tctgagggga ctagggtgtg 7080
tttaggcgaa aagcggggct tcggttgtac gcggttagga gtccccttag gatatagtag 7140
tttcgctttt gcatagggag ggggaaatgt agtcttatgc aatactcttg tagtcttgca 7200
acatggtaac gatgagttag caacatgcct tacaaggaga gaaaaagcac cgtgcatgcc 7260
gattggtgga agtaagggtg tacgatectg ccttattagg aaggcaacag acgggtctga 7320
catggattgg acgaaccact gaattccgca ttgcagagat attgtattta agtgcctagc 7380
tcgatacaat aaacgccatt tgaccattca ccacattggg gtgcacctgg gttgatggcc 7440
ggaccgttga ttccctgacg actacgagca cctgcatgaa gcagaaggct tcatttggtg 7500
accccgacgt gatagttagg gaatagtggg cggccacaga cggcgtggcg atcctgtctc 7560
catcctctc gtctatcggg aggcgacttc gatgacctg gtggaggggg ctgcggctta 7620
gggaggcaga agctgagtac cgtcggaggg gatccacagg acgggtgtgg tcgccatgat 7680
cgcgtagtgc atagtggctc caagtagcga agcagcagg actgggcggc ggccaaagcg 7740
gtcggacagt gctccgagaa cgggtgcgca tagaaattgc atcaacgcat atagcgctag 7800
cagcacgcca tagtgactgg cgatgctgtc ggaatggacg atatcccga agaggccccg 7860
cagtaccggc ataaccaagc ctatgcctac agcatccagg gtgacggtgc cgaggatgac 7920
gatgagcgca ttgttagatt tcatacacgg tgcctgactg cgtttagcaat ttaactgtga 7980
taaactaccg cattaaagct ccaaacttgg ctgtttcctg tgtgaaattg ttatccgctc 8040
acaattccac acattatacg agccggaagc ataaagtgtg aaacctgggg tgcctaataga 8100
gtgagaattc ttgaagacga aagggcctcg tgatacgctt atttttatag gttaatgtca 8160

9/20

tgataataat ggtttcttag acgtcaggtg gcacttttcg gggaaatgtg cgcggaaccc 8220
ctatttgttt atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct 8280
gataaatgct tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg 8340
cccttattcc cttttttgcg gcattttgccc ttctgtttt tgctcaccga gaaacgctgg 8400
tgaaagtaaa agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc 8460
tcaacagcgg taagatcctt gagagttttc gccccgaaga acgtttttcca atgatgagca 8520
cttttaaagt tctgctatgt ggcgcggtat tatcccggtg tgacgcggg caagagcaac 8580
tcggtcgccc catacactat tctcagaatg acttggttga gtactacca gtcacagaaa 8640
agcatcttac ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg 8700
ataacactgc ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt 8760
ttttgcacaa catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg 8820
aagccatacc aaacgacgag cgtgacacca cgatgcctgc agcaatggca acaacgttgc 8880
gcaaaactatt aactggcgaa ctacttactc tagcttcccg gcaacaatta atagactgga 8940
tgaggcgga taaagttgca ggaccacttc tgcgctcggc cttccggct ggctggttta 9000
ttgctgataa atctggagcc ggtgagcgtg ggtctcgcg tatcattgca gcactggggc 9060
cagatggtaa gccctcccg atcgtagtta tctacacgac ggggagtcag gcaactatgg 9120
atgaacgaaa tagacagatc gctgagatag gtgcctcact gattaagcat tggtaactgt 9180
cagaccaagt ttactcatat atactttaga ttgatttaa acttcatttt taatttaaaa 9240
ggatctaggt gaagatcctt tttgataatc tcatgaccaa aatcccttaa cgtgagtttt 9300
cgttccactg agcgtcagac cccgtagaaa agatcaaagg atcttcttga gatccttttt 9360
ttctgcgct aatctgctgc ttgcaaacaa aaaaaccacc gctaccagcg gtggtttgtt 9420
tgccggatca agagctacca actctttttc cgaaggtaac tggcttcagc agagcgaga 9480
taccaaatac tgtccttcta gtgtagccgt agttaggcca ccacttcaag aactctgtag 9540
caccgcctac atacctcgct ctgctaatec tgttaccagt ggctgctgcc agtggcgata 9600
agtcgtgtct taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg 9660
gctgaacggg gggttcgtgc acacagccca gcttgagcgc aacgacctac accgaactga 9720
gatacctaca gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcgga 9780
ggatatccgt aagcggcagg gtcggaacag gagagcgcac gagggagctt ccagggggaa 9840
acgcctggt tctttatagt cctgtcgggt ttcccaacct ctgacttgag cgtcgatttt 9900
tgtgatgctc gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac 9960

10/20

ggttcctggc cttttgctgg ccttttgctc acatgttctt tectgcgta tcccttgatt 10020
ctgtggataa ccgtattacc gcctttgagt gagctgatac cgctcgccgc agccgaacga 10080
ccgagcgcag cgagtcagtg agcgaggaag cggaagagcg cctgatgcgg tattttctcc 10140
ttacgcatct gtgcggtatt tcacaccgca tatgggtgcac tctcagtaca atctgctctg 10200
atgccgcata gttaagccag tatacactcc gctatcgcta cgtgactggg tcatggctgc 10260
gccccgacac ccgccaacac ccgctgacgc gccctgacgg gcttgtctgc tcccggcatc 10320
cgcttacaga caagctgtga ccgtctccgg gagctgcatg tgtcagaggt ttccaccgtc 10380
atcaccgaaa cgcgcgaggg agctgcggta aagctcatca gcgtggctgt gaagcgattc 10440
acagatgtct gcctgttcat ccgcgtccag ctcgttgagt ttctccagaa gcgttaatgt 10500
ctggcttctg ataaagcggg ccatgttaag ggcggttttt tctgttttg tcaattgatg 10560
cctcgtgta agggggaatt tctgttcatg ggggtaatga taccgatgaa acgagagagg 10620
atgctcacga tacgggttac tgatgatgaa catgcccggg tactggaacg ttgtgagggg 10680
aaacaactgg cggatatgat gcggcgggac cagagaaaaa tcaactcaggg tcaatgccag 10740
cgcttcgtta atacagatgt aggtgttcca cagggtagcc agcagcatcc tgcgatgcag 10800
atccggaaca taatggtgca gggcgtgac ttccgcgttt ccagacttta cgaaacacgg 10860
aaaccgaaga ccattcatgt tgttgctcag gtcgcagacg ttttgacga gcagtcgctt 10920
cacgttcgct cgcgtatcgg tgattcatc tgctaaccag taaggcaacc ccgccagcct 10980
agccgggtcc tcaacgacag gagcacgac atgagcacc gtggccagga cccaacgctg 11040
cccagatgc gccgcgtgcg gcgctggag atggcggacg cgatggatat gttctgccaa 11100
gggttggttt gcgcattcac agttctccgc aagaattgat tggctccaat tcttgagtg 11160
gtgaatccgt tagcgaggtg ccgcgggctt ccattcaggt cgaggtggcc cggctccatg 11220
caccgcgacg caacgcgggg aggcagacaa ggtatagggc ggcgatgca tgtacgggcc 11280
agatatacgc gtatctgagg ggactaggg gtgttttaggc gaaaagcggg gcttcggttg 11340
tacgcggtta ggagtccct taggatatag tagtttcgct tttgcatagg gagggggaaa 11400
tgtagtetta tgcaatactc ttgtagtctt gcaacatggt aacgatgagt tagcaacatg 11460
ccttacaagg agagaaaaag caccgtgcat gccgattggg ggaagtaagg tggtagcatc 11520
gtgccttatt aggaaggcaa cagacgggtc tgacatggat tggacgaacc actgaattcc 11580
gcattgcaga gatattgtat ttaagtgcct agctcgatac aataaac 11627

11/20

<210> 2
 <211> 1759
 <212> DNA
 <213> Chicken

<220>

<221> gene
 <222> (1)..(1759)
 <223> chicken c-SRC cDNA

<220>

<221> CDS
 <222> (112)..(1710)

<400> 2
 tctgacaccc atctgtctgt ctgtctgtgt gctgcaggag ctgagctgac tctgctgtgg 60
 cctogcgtac cactgtggcc aggcggtagc tgggacgtgc agcccaccac c atg ggg 117
 Met Gly
 1
 agc agc aag agc aag ccc aag gac ccc agc cag cgc cgg cgc agc ctg 165
 Ser Ser Lys Ser Lys Pro Lys Asp Pro Ser Gln Arg Arg Arg Ser Leu
 5 10 15
 gag cca ccc gac agc acc cac cac ggg gga ttc cca gcc tcg cag acc 213
 Glu Pro Pro Asp Ser Thr His His Gly Gly Phe Pro Ala Ser Gln Thr
 20 25 30
 ccc aac aag aca gca gcc ccc gac acg cac cgc acc ccc agc cgc tcc 261
 Pro Asn Lys Thr Ala Ala Pro Asp Thr His Arg Thr Pro Ser Arg Ser
 35 40 45 50
 ttt ggg acc gtg gcc acc gag ccc aag ctc ttc ggg ggc ttc aac act 309
 Phe Gly Thr Val Ala Thr Glu Pro Lys Leu Phe Gly Gly Phe Asn Thr
 55 60 65
 tct gac acc gtt acg tcg ccg cag cgt gcc ggg gca ctg gct ggc ggc 357
 Ser Asp Thr Val Thr Ser Pro Gln Arg Ala Gly Ala Leu Ala Gly Gly
 70 75 80
 gtc acc act ttc gtg gct ctc tac gac tac gag tcc cgg act gaa acg 405
 Val Thr Thr Phe Val Ala Leu Tyr Asp Tyr Glu Ser Arg Thr Glu Thr
 85 90 95
 gac ttg tcc ttc aag aaa gga gaa cgc ctg cag att gtc aac aac acg 453
 Asp Leu Ser Phe Lys Lys Gly Glu Arg Leu Gln Ile Val Asn Asn Thr
 100 105 110
 gaa ggt gac tgg tgg ctg gct cat tcc ctc act aca gga cag acg ggc 501
 Glu Gly Asp Trp Trp Leu Ala His Ser Leu Thr Thr Gly Gln Thr Gly
 115 120 125 130

12/20

| | |
|---|------|
| tac atc ccc agt aac tat gtc gcg ccc tca gac tcc atc cag gct gaa | 549 |
| Tyr Ile Pro Ser Asn Tyr Val Ala Pro Ser Asp Ser Ile Gln Ala Glu | |
| 135 140 145 | |
| gag tgg tac ttt ggg aag atc act cgt cgg gag tcc gag cgg ctg ctg | 597 |
| Glu Trp Tyr Phe Gly Lys Ile Thr Arg Arg Glu Ser Glu Arg Leu Leu | |
| 150 155 160 | |
| ctc aac ccc gaa aac ccc cgg gga acc ttc ttg gtc cgg gag agc gag | 645 |
| Leu Asn Pro Glu Asn Pro Arg Gly Thr Phe Leu Val Arg Glu Ser Glu | |
| 165 170 175 | |
| acg aca aaa ggt gcc tat tgc ctc tcc gtt tct gac ttt gac aac gcc | 693 |
| Thr Thr Lys Gly Ala Tyr Cys Leu Ser Val Ser Asp Phe Asp Asn Ala | |
| 180 185 190 | |
| aag ggg ctc aat gtg aag cac tac aag atc cgc aag ctg gac agc ggc | 741 |
| Lys Gly Leu Asn Val Lys His Tyr Lys Ile Arg Lys Leu Asp Ser Gly | |
| 195 200 205 210 | |
| ggc ttc tac atc acc tca cgc aca cag ttc agc agc ctg cag cag ctg | 789 |
| Gly Phe Tyr Ile Thr Ser Arg Thr Gln Phe Ser Ser Leu Gln Gln Leu | |
| 215 220 225 | |
| gtg gcc tac tac tcc aaa cat gct gat ggc ttg tgc cac cgc ctg acc | 837 |
| Val Ala Tyr Tyr Ser Lys His Ala Asp Gly Leu Cys His Arg Leu Thr | |
| 230 235 240 | |
| aac gtc tgc ccc acg tcc aag ccc cag acc cag gga ctc gcc aag gac | 885 |
| Asn Val Cys Pro Thr Ser Lys Pro Gln Thr Gln Gly Leu Ala Lys Asp | |
| 245 250 255 | |
| gcg tgg gaa atc ccc cgg gag tcg ctg cgg ctg gag gtg aag ctg ggg | 933 |
| Ala Trp Glu Ile Pro Arg Glu Ser Leu Arg Leu Glu Val Lys Leu Gly | |
| 260 265 270 | |
| cag ggc tgc ttt gga gag gtc tgg atg ggg acc tgg aac ggc acc acc | 981 |
| Gln Gly Cys Phe Gly Glu Val Trp Met Gly Thr Trp Asn Gly Thr Thr | |
| 275 280 285 290 | |
| aga gtg gcc ata aag act ctg aag ccc ggc acc atg tcc ccg gag gcc | 1029 |
| Arg Val Ala Ile Lys Thr Leu Lys Pro Gly Thr Met Ser Pro Glu Ala | |
| 295 300 305 | |
| ttc ctg cag gaa gcc caa gtg atg aag aag ctc cgg cat gag aag ctg | 1077 |
| Phe Leu Gln Glu Ala Gln Val Met Lys Lys Leu Arg His Glu Lys Leu | |
| 310 315 320 | |
| gtt cag ctg tac gca gtg gtg tcg gaa gag ccc atc tac atc gtc act | 1125 |
| Val Gln Leu Tyr Ala Val Val Ser Glu Glu Pro Ile Tyr Ile Val Thr | |
| 325 330 335 | |
| gag tac atg agc aag ggg agc ctc ctg gat ttc ctg aag gga gag atg | 1173 |
| Glu Tyr Met Ser Lys Gly Ser Leu Leu Asp Phe Leu Lys Gly Glu Met | |
| 340 345 350 | |
| ggc aag tac ctg cgg ctg cca cag ctc gtc gat atg gct gct cag att | 1221 |
| Gly Lys Tyr Leu Arg Leu Pro Gln Leu Val Asp Met Ala Ala Gln Ile | |
| 355 360 365 370 | |

14/20

Gln Thr Pro Asn Lys Thr Ala Ala Pro Asp Thr His Arg Thr Pro Ser
 35 40 45
 Arg Ser Phe Gly Thr Val Ala Thr Glu Pro Lys Leu Phe Gly Gly Phe
 50 55 60
 Asn Thr Ser Asp Thr Val Thr Ser Pro Gln Arg Ala Gly Ala Leu Ala
 65 70 75 80
 Gly Gly Val Thr Thr Phe Val Ala Leu Tyr Asp Tyr Glu Ser Arg Thr
 85 90 95
 Glu Thr Asp Leu Ser Phe Lys Lys Gly Glu Arg Leu Gln Ile Val Asn
 100 105 110
 Asn Thr Glu Gly Asp Trp Trp Leu Ala His Ser Leu Thr Thr Gly Gln
 115 120 125
 Thr Gly Tyr Ile Pro Ser Asn Tyr Val Ala Pro Ser Asp Ser Ile Gln
 130 135 140
 Ala Glu Glu Trp Tyr Phe Gly Lys Ile Thr Arg Arg Glu Ser Glu Arg
 145 150 155 160
 Leu Leu Leu Asn Pro Glu Asn Pro Arg Gly Thr Phe Leu Val Arg Glu
 165 170 175
 Ser Glu Thr Thr Lys Gly Ala Tyr Cys Leu Ser Val Ser Asp Phe Asp
 180 185 190
 Asn Ala Lys Gly Leu Asn Val Lys His Tyr Lys Ile Arg Lys Leu Asp
 195 200 205
 Ser Gly Gly Phe Tyr Ile Thr Ser Arg Thr Gln Phe Ser Ser Leu Gln
 210 215 220
 Gln Leu Val Ala Tyr Tyr Ser Lys His Ala Asp Gly Leu Cys His Arg
 225 230 235 240
 Leu Thr Asn Val Cys Pro Thr Ser Lys Pro Gln Thr Gln Gly Leu Ala
 245 250 255
 Lys Asp Ala Trp Glu Ile Pro Arg Glu Ser Leu Arg Leu Glu Val Lys
 260 265 270
 Leu Gly Gln Gly Cys Phe Gly Glu Val Trp Met Gly Thr Trp Asn Gly
 275 280 285
 Thr Thr Arg Val Ala Ile Lys Thr Leu Lys Pro Gly Thr Met Ser Pro
 290 295 300
 Glu Ala Phe Leu Gln Glu Ala Gln Val Met Lys Lys Leu Arg His Glu
 305 310 315 320
 Lys Leu Val Gln Leu Tyr Ala Val Val Ser Glu Glu Pro Ile Tyr Ile
 325 330 335

15/20

Val Thr Glu Tyr Met Ser Lys Gly Ser Leu Leu Asp Phe Leu Lys Gly
 340 345 350
 Glu Met Gly Lys Tyr Leu Arg Leu Pro Gln Leu Val Asp Met Ala Ala
 355 360 365
 Gln Ile Ala Ser Gly Met Ala Tyr Val Glu Arg Met Asn Tyr Val His
 370 375 380
 Arg Asp Leu Arg Ala Ala Asn Ile Leu Val Gly Glu Asn Leu Val Cys
 385 390 395 400
 Lys Val Ala Asp Phe Gly Leu Ala Arg Leu Ile Glu Asp Asn Glu Tyr
 405 410 415
 Thr Ala Arg Gln Gly Ala Lys Phe Pro Ile Lys Trp Thr Ala Pro Glu
 420 425 430
 Ala Ala Leu Tyr Gly Arg Phe Thr Ile Lys Ser Asp Val Trp Ser Phe
 435 440 445
 Gly Ile Leu Leu Thr Glu Leu Thr Thr Lys Gly Arg Val Pro Tyr Pro
 450 455 460
 Gly Met Val Asn Arg Glu Val Leu Asp Gln Val Glu Arg Gly Tyr Arg
 465 470 475 480
 Met Pro Cys Pro Pro Glu Cys Pro Glu Ser Leu His Asp Leu Met Cys
 485 490 495
 Gln Cys Trp Arg Arg Asp Pro Glu Glu Arg Pro Thr Phe Glu Tyr Leu
 500 505 510
 Gln Ala Phe Leu Glu Asp Tyr Phe Thr Ser Thr Glu Pro Gln Tyr Gln
 515 520 525
 Pro Gly Glu Asn Leu
 530

<210> 4
 <211> 2187
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> gene
 <222> (1)..(2187)
 <223> human c-SRC cDNA

<220>
 <221> CDS
 <222> (134)..(1483)

16/20

<400> 4

gcgcgcgcgc cgcagggccg tgatgccgcc cgcgcggagg tggcccggac cgcagtgccc 60
 caagagagct ctaatggtac caagtgcacg gttggcttta ctgtgactcg gggacgccag 120
 agctcctgag aag atg tca gca ata cag gcc gcc tgg cca tcc ggt aca 169
 Met Ser Ala Ile Gln Ala Ala Trp Pro Ser Gly Thr
 1 5 10
 gaa tgt att gcc aag tac aac ttc cac ggc act gcc gag cag gac ctg 217
 Glu Cys Ile Ala Lys Tyr Asn Phe His Gly Thr Ala Glu Gln Asp Leu
 15 20 25
 ccc ttc tgc aaa gga gac gtg ctc acc att gtg gcc gtc acc aag gac 265
 Pro Phe Cys Lys Gly Asp Val Leu Thr Ile Val Ala Val Thr Lys Asp
 30 35 40
 ccc aac tgg tac aaa gcc aaa aac aag gtg ggc cgt gag ggc atc atc 313
 Pro Asn Trp Tyr Lys Ala Lys Asn Lys Val Gly Arg Glu Gly Ile Ile
 45 50 55 60
 cca gcc aac tac gtc cag aag cgg gag ggc gtg aag gcg ggt acc aaa 361
 Pro Ala Asn Tyr Val Gln Lys Arg Glu Gly Val Lys Ala Gly Thr Lys
 65 70 75
 ctc agc ctc atg cct tgg ttc cac ggc aag atc aca cgg gag cag gct 409
 Leu Ser Leu Met Pro Trp Phe His Gly Lys Ile Thr Arg Glu Gln Ala
 80 85 90
 gag cgg ctt ctg tac ccg ccg gag aca ggc ctg ttc ctg gtg cgg gag 457
 Glu Arg Leu Leu Tyr Pro Pro Glu Thr Gly Leu Phe Leu Val Arg Glu
 95 100 105
 agc acc aac tac ccc gga gac tac acg ctg tgc gtg agc tgc gac ggc 505
 Ser Thr Asn Tyr Pro Gly Asp Tyr Thr Leu Cys Val Ser Cys Asp Gly
 110 115 120
 aag gtg gag cac tac cgc atc atg tac cat gcc agc aag ctc agc atc 553
 Lys Val Glu His Tyr Arg Ile Met Tyr His Ala Ser Lys Leu Ser Ile
 125 130 135 140
 gac gag gag gtg tac ttt gag aac ctc atg cag ctg gtg gag cac tac 601
 Asp Glu Glu Val Tyr Phe Glu Asn Leu Met Gln Leu Val Glu His Tyr
 145 150 155
 acc tca gac gca gat gga ctc tgt acg cgc ctc att aaa cca aag gtc 649
 Thr Ser Asp Ala Asp Gly Leu Cys Thr Arg Leu Ile Lys Pro Lys Val
 160 165 170
 atg gag ggc aca gtg gcg gcc cag gat gag ttc tac cgc agc ggc tgg 697
 Met Glu Gly Thr Val Ala Ala Gln Asp Glu Phe Tyr Arg Ser Gly Trp
 175 180 185
 gcc ctg aac atg aag gag ctg aag ctg ctg cag acc atc ggg aag ggg 745
 Ala Leu Asn Met Lys Glu Leu Lys Leu Leu Gln Thr Ile Gly Lys Gly
 190 195 200

17/20

| | |
|---|------|
| gag ttc gga gac gtg atg ctg ggc gat tac cga ggg aac aaa gtc gcc | 793 |
| Glu Phe Gly Asp Val Met Leu Gly Asp Tyr Arg Gly Asn Lys Val Ala | |
| 205 210 215 220 | |
| gtc aag tgc att aag aac gac gcc act gcc cag gcc ttc ctg gct gaa | 841 |
| Val Lys Cys Ile Lys Asn Asp Ala Thr Ala Gln Ala Phe Leu Ala Glu | |
| 225 230 235 | |
| gcc tca gtc atg acg caa ctg cgg cat agc aac ctg gtg cag ctc ctg | 889 |
| Ala Ser Val Met Thr Gln Leu Arg His Ser Asn Leu Val Gln Leu Leu | |
| 240 245 250 | |
| ggc gtg atc gtg gag gag aag ggc ggg ctc tac atc gtc act gag tac | 937 |
| Gly Val Ile Val Glu Glu Lys Gly Gly Leu Tyr Ile Val Thr Glu Tyr | |
| 255 260 265 | |
| atg gcc aag ggg agc ctt gtg gac tac ctg cgg tct agg ggt cgg tca | 985 |
| Met Ala Lys Gly Ser Leu Val Asp Tyr Leu Arg Ser Arg Gly Arg Ser | |
| 270 275 280 | |
| gtg ctg ggc gga gac tgt ctc ctc aag ttc tgc cta gat gtc tgc gag | 1033 |
| Val Leu Gly Gly Asp Cys Leu Leu Lys Phe Ser Leu Asp Val Cys Glu | |
| 285 290 295 300 | |
| gcc atg gaa tac ctg gag ggc aac aat ttc gtg cat cga gac ctg gct | 1081 |
| Ala Met Glu Tyr Leu Glu Gly Asn Asn Phe Val His Arg Asp Leu Ala | |
| 305 310 315 | |
| gcc cgc aat gtg ctg gtg tct gag gac aac gtg gcc aag gtc agc gac | 1129 |
| Ala Arg Asn Val Leu Val Ser Glu Asp Asn Val Ala Lys Val Ser Asp | |
| 320 325 330 | |
| ttt ggt ctc acc aag gag gcg tcc agc acc cag gac acg ggc aag ctg | 1177 |
| Phe Gly Leu Thr Lys Glu Ala Ser Ser Thr Gln Asp Thr Gly Lys Leu | |
| 335 340 345 | |
| cca gtc aag tgg aca gcc cct gag gcc ctg aga gag aag aaa ttc tcc | 1225 |
| Pro Val Lys Trp Thr Ala Pro Glu Ala Leu Arg Glu Lys Lys Phe Ser | |
| 350 355 360 | |
| act aag tct gac gtg tgg agt ttc gga atc ctt ctc tgg gaa atc tac | 1273 |
| Thr Lys Ser Asp Val Trp Ser Phe Gly Ile Leu Leu Trp Glu Ile Tyr | |
| 365 370 375 380 | |
| tcc ttt ggg cga gtg cct tat cca aga att ccc ctg aag gac gtc gtc | 1321 |
| Ser Phe Gly Arg Val Pro Tyr Pro Arg Ile Pro Leu Lys Asp Val Val | |
| 385 390 395 | |
| cct cgg gtg gag aag ggc tac aag atg gat gcc ccc gac ggc tgc ccg | 1369 |
| Pro Arg Val Glu Lys Gly Tyr Lys Met Asp Ala Pro Asp Gly Cys Pro | |
| 400 405 410 | |
| ccc gca gtc tat gaa gtc atg aag aac tgc tgg cac ctg gac gcc gcc | 1417 |
| Pro Ala Val Tyr Glu Val Met Lys Asn Cys Trp His Leu Asp Ala Ala | |
| 415 420 425 | |

18/20

atg cgg ccc tcc ttc cta cag ctc cga gag cag ctt gag cac atc aaa 1465
 Met Arg Pro Ser Phe Leu Gln Leu Arg Glu Gln Leu Glu His Ile Lys
 430 435 440

acc cac gag ctg cac ctg tgacggctgg cctccgctg ggtcatgggc 1513
 Thr His Glu Leu His Leu
 445 450

ctgtggggac tgaacctgga agatcatgga cctgggtgcc ctgctcactg ggcccagacc 1573
 tgaactgagc cccagcgggc tggcgggcct ttttctctgcg tcccagcctg caccctccg 1633
 gccccgtctc tcttggaacc acctgtgggg cctggggagc cactgaggg gccagggagg 1693
 aaggaggcca cggagcggga ggcagcgccc caccacgtcg ggcttccctg gcctcccgcc 1753
 actcgcttc ttagagtttt attcctttcc ttttttgaga tttttttcc gtgtgtttat 1813
 tttttattat ttttcaagat aaggagaaag aaagtaccca gcaaattggc attttacaag 1873
 aagtacgaat cttatttttc ctgtctctgc cgtgaggggtg ggggggaccg ggccctctc 1933
 tagggacccc tcgccccagc ctcattcccc attctgtgtc ccatgtcccg tgtctctcg 1993
 gtgcgccgt gtttgcgctt gaccatgttg cactgtttgc atgcgcccga ggcagacgtc 2053
 tgtcaggggc ttggatttcg tgtgccgtg ccaccgccc acccgcttg tgagatggaa 2113
 ttgtaataaa ccacgccatg aggacaccgc cgccgcctc ggcgcttct ccaccgaaaa 2173
 aaaaaaaaaa aaaa 2187

<210> 5
 <211> 450
 <212> PRT
 <213> Homo sapiens

<400> 5
 Met Ser Ala Ile Gln Ala Ala Trp Pro Ser Gly Thr Glu Cys Ile Ala
 1 5 10 15
 Lys Tyr Asn Phe His Gly Thr Ala Glu Gln Asp Leu Pro Phe Cys Lys
 20 25 30
 Gly Asp Val Leu Thr Ile Val Ala Val Thr Lys Asp Pro Asn Trp Tyr
 35 40 45
 Lys Ala Lys Asn Lys Val Gly Arg Glu Gly Ile Ile Pro Ala Asn Tyr
 50 55 60
 Val Gln Lys Arg Glu Gly Val Lys Ala Gly Thr Lys Leu Ser Leu Met
 65 70 75 80

19/20

| | | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|
| Pro | Trp | Phe | His | Gly | Lys | Ile | Thr | Arg | Glu | Gln | Ala | Glu | Arg | Leu | Leu | | |
| | | | | 85 | | | | | 90 | | | | | 95 | | | |
| Tyr | Pro | Pro | Glu | Thr | Gly | Leu | Phe | Leu | Val | Arg | Glu | Ser | Thr | Asn | Tyr | | |
| | | | 100 | | | | | 105 | | | | | 110 | | | | |
| Pro | Gly | Asp | Tyr | Thr | Leu | Cys | Val | Ser | Cys | Asp | Gly | Lys | Val | Glu | His | | |
| | | 115 | | | | | 120 | | | | | 125 | | | | | |
| Tyr | Arg | Ile | Met | Tyr | His | Ala | Ser | Lys | Leu | Ser | Ile | Asp | Glu | Glu | Val | | |
| | 130 | | | | | 135 | | | | | 140 | | | | | | |
| Tyr | Phe | Glu | Asn | Leu | Met | Gln | Leu | Val | Glu | His | Tyr | Thr | Ser | Asp | Ala | | |
| | 145 | | | | 150 | | | | | 155 | | | | | 160 | | |
| Asp | Gly | Leu | Cys | Thr | Arg | Leu | Ile | Lys | Pro | Lys | Val | Met | Glu | Gly | Thr | | |
| | | | | 165 | | | | | 170 | | | | | 175 | | | |
| Val | Ala | Ala | Gln | Asp | Glu | Phe | Tyr | Arg | Ser | Gly | Trp | Ala | Leu | Asn | Met | | |
| | | | 180 | | | | | 185 | | | | | 190 | | | | |
| Lys | Glu | Leu | Lys | Leu | Leu | Gln | Thr | Ile | Gly | Lys | Gly | Glu | Phe | Gly | Asp | | |
| | | 195 | | | | | 200 | | | | | 205 | | | | | |
| Val | Met | Leu | Gly | Asp | Tyr | Arg | Gly | Asn | Lys | Val | Ala | Val | Lys | Cys | Ile | | |
| | 210 | | | | | 215 | | | | | 220 | | | | | | |
| Lys | Asn | Asp | Ala | Thr | Ala | Gln | Ala | Phe | Leu | Ala | Glu | Ala | Ser | Val | Met | | |
| | 225 | | | | 230 | | | | | 235 | | | | | 240 | | |
| Thr | Gln | Leu | Arg | His | Ser | Asn | Leu | Val | Gln | Leu | Leu | Gly | Val | Ile | Val | | |
| | | | | 245 | | | | | 250 | | | | | 255 | | | |
| Glu | Glu | Lys | Gly | Gly | Leu | Tyr | Ile | Val | Thr | Glu | Tyr | Met | Ala | Lys | Gly | | |
| | | | 260 | | | | | 265 | | | | | 270 | | | | |
| Ser | Leu | Val | Asp | Tyr | Leu | Arg | Ser | Arg | Gly | Arg | Ser | Val | Leu | Gly | Gly | | |
| | | 275 | | | | | 280 | | | | | 285 | | | | | |
| Asp | Cys | Leu | Leu | Lys | Phe | Ser | Leu | Asp | Val | Cys | Glu | Ala | Met | Glu | Tyr | | |
| | 290 | | | | | 295 | | | | | 300 | | | | | | |
| Leu | Glu | Gly | Asn | Asn | Phe | Val | His | Arg | Asp | Leu | Ala | Ala | Arg | Asn | Val | | |
| | 305 | | | | 310 | | | | | 315 | | | | | 320 | | |
| Leu | Val | Ser | Glu | Asp | Asn | Val | Ala | Lys | Val | Ser | Asp | Phe | Gly | Leu | Thr | | |
| | | | | 325 | | | | | 330 | | | | | 335 | | | |
| Lys | Glu | Ala | Ser | Ser | Thr | Gln | Asp | Thr | Gly | Lys | Leu | Pro | Val | Lys | Trp | | |
| | | | 340 | | | | | 345 | | | | | 350 | | | | |
| Thr | Ala | Pro | Glu | Ala | Leu | Arg | Glu | Lys | Lys | Phe | Ser | Thr | Lys | Ser | Asp | | |
| | | 355 | | | | | 360 | | | | | 365 | | | | | |
| Val | Trp | Ser | Phe | Gly | Ile | Leu | Leu | Trp | Glu | Ile | Tyr | Ser | Phe | Gly | Arg | | |
| | 370 | | | | | 375 | | | | | 380 | | | | | | |

20/20

Val Pro Tyr Pro Arg Ile Pro Leu Lys Asp Val Val Pro Arg Val Glu
 385 390 395 400

Lys Gly Tyr Lys Met Asp Ala Pro Asp Gly Cys Pro Pro Ala Val Tyr
 405 410 415

Glu Val Met Lys Asn Cys Trp His Leu Asp Ala Ala Met Arg Pro Ser
 420 425 430

Phe Leu Gln Leu Arg Glu Gln Leu Glu His Ile Lys Thr His Glu Leu
 435 440 445

His Leu
 450

<210> 6

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:9E10-myc
 epitope tag

<400> 6

Val Asp Met Glu Gln Lys Leu Ile Ala Glu Glu Asp Leu Asn
 1 5 10

WO 99/61590

PCT/US99/11780

1/18

SEQUENCE LISTING

<110> The Scripps Research Institute et al.

<120> METHODS AND COMPOSITIONS USEFUL FOR MODULATION OF
ANGIOGENESIS USING TYROSINE KINASE SRC

<130> TSRI 651.1

<140> Not yet known
<141> To be determined

<150> 60/087,220
<151> 1998-05-29

<160> 6

<170> PatentIn Ver. 2.0

<210> 1
<211> 11627
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:RCASBP(A) based
on avian sarcoma virus

<220>
<221> misc_feature
<222> (7649)..(11258)
<223> pBR322 sequences

<220>
<221> LTR
<222> (7166)..(7494)
<223> upstream

<220>
<221> LTR
<222> (1)..(101)
<223> upstream (numbering begins at the upstream R)

<220>
<221> misc_feature
<222> (11394)..(11623)
<223> U3

<220>
<221> misc_feature
<222> (1)..(21)
<223> R

<220>
<221> misc_feature
<222> (22)..(101)
<223> U5

<220>
<221> misc_feature
<222> (102)..(119)

<220>
<221> LTR

WO 99/61590

PCT/US99/11780

2/18

<222> (7166)..(7494)
<223> downstream

<220>
<221> misc_feature
<222> (7166)..(7393)
<223> U3

<220>
<221> misc_feature
<222> (7394)..(7414)
<223> R

<220>
<221> misc_feature
<222> (7415)..(7494)
<223> U5

<220>
<221> misc_feature
<222> (7154)..(7165)
<223> PPT

<220>
<221> misc_feature
<222> (388)..(391)
<223> splice donor (AGGT)

<220>
<221> misc_feature
<222> (5074)..(5077)
<223> env splice acceptor (AGGC)

<220>
<221> misc_feature
<222> (6982)..(6985)
<223> ClaI splice acceptor (AGGA)

<220>
<221> gene
<222> (372)..(902)
<223> gag p19

<220>
<221> gene
<222> (909)..(1094)
<223> gag p10

<220>
<221> gene
<222> (1095)..(1814)
<223> gag p27

<220>
<221> gene
<222> (1843)..(2108)
<223> gag p12

<220>
<221> gene
<222> (2109)..(2480)
<223> gag p15

<220>

WO 99/61590

PCT/US99/11780

3/18

<221> misc_signal
 <222> (2481)..(2483)
 <223> gag stop

<220>
 <221> gene
 <222> (2501)..(4216)
 <223> pol RT

<220>
 <221> gene
 <222> (4217)..(5185)
 <223> pol IN

<220>
 <221> misc_signal
 <222> (5186)..(5188)
 <223> pol stop

<220>
 <221> gene
 <222> (5244)..(6263)
 <223> env gp85

<220>
 <221> gene
 <222> (6264)..(6878)
 <223> env gp37

<220>
 <221> misc_signal
 <222> (6879)..(6881)
 <223> env stop

<220>
 <221> misc_feature
 <222> (7027)
 <223> ClaI site/ the ClaI site in gag is methylated in
 Dam+ strains and does not cut.

<400> 1
 gccatttgac cattcaccac attggtgtgc acctgggttg atggccggac cgttgattcc 60
 ctgacgacta cgagcacctg catgaagcag aaggtttcat ttggtgaccc cgacgtgata 120
 gttaggggaat agtggtcggc cacagacggc gtggcgatcc tgtctccatc cgtctcgtct 180
 atcgggaggg gagttcgatg accctgggtg agggggctgc ggcttaggga ggcagaagct 240
 gagtaccgtc ggaggggagct ccagggcccc gagcgactga cccctgccga gaactcagag 300
 ggtcgtcgga agacggagag tgagcccgac gaccacccca ggcacgtctt tggtcggcct 360
 gcggatcaag catggaagcc gtcattaagg tgatttcgtc cgcgtgtaaa acctattgcy 420
 ggaaaatctc tccttctaag aaggaaatcg gggccatgtt gtccctgtta caaaaaggaa 480
 gggttgcttat gtctccctca gatattatatt ctccgggggtc ctgggatccc atcactgcgg 540
 cgctctccca gcgggcaatg gtacttgga aatcggggaga gttaaaaacc tggggattgg 600
 ttttgggggc attgaaggcg gctcgagagg aacagggttac atctgagcaa gcaaagtgtt 660
 gggtgggatt aggggggagg aggggtctct ccccagggtc ggagtgcac gagaaaccag 720

ctacggagcg gcgaatcgac aaaggggagg aggtggggaga aacaactgtg cagcgagatg 780
cgaagatggc gccagaggaa gcggccacac ctaaaaccgt tggcacatcc tgctatcatt 840
gcggaacagc tgttggctgc aattgcgcca ccgccacagc ctcggeccct cctccccctt 900
atgtggggag tggtttgtat ccttccctgg cgggggtggg agagcagcag ggccagggag 960
ataacacgtc tcggggggcg gagcagccaa gggaggagcc agggcacgcg ggtcaggccc 1020
ctgggcccgc cctgactgac tgggcaaggg taaggaggga gcttgcgagt actggtccgc 1080
ccgtggtggc catgcctgta gtgattaaga cagagggacc cgcctggacc cctctggagc 1140
caaaattgat cacaagactg gctgatacgg tcaggaccaa gggettacga tccccgatca 1200
ctatggcaga agtggaagcg ctcatgtcct ccccgttgct gccgcatgac gtcacgaatc 1260
taatgagagt gatttttaga cctgccccat atgccttatg gatggacgct tggggagtc 1320
aactccagac ggttatagcg gcagccactc gcgacccccg acaccagcg aacggtcaag 1380
ggcgggggga acggactaac ttggatcgat taaagggtt agctgatggg atggtgggca 1440
accacaggg tcaggccgca ttattaagac cgggggaatt ggttgctatt acggcgtcgg 1500
ctctccaggc gtttagagaa gttgcccggc tggcggaacc tgcaggtcca tggcgggaca 1560
tcacgcagg accatctgag tcctttgttg attttgccaa tcggcttata aaggcggttg 1620
aggggtcaga tctcccgct tccgcgcggg ctccggtgat cattgactgc ttaggcaga 1680
agtcacagcc agatattcag cagcttatac gggcagcacc ctccacgctg accacccag 1740
gagagataat caaatatgtg ctagacaggc agaagattgc ccctcttacg gatcaaggca 1800
tagccgccc catgtcgtct gctatccagc ccttagttat ggcagtagtc aatagagaga 1860
gggatggaca aactgggtcg ggtggtcgtg cccgagggt ctgctacact tgtggatccc 1920
cgggacatta tcaggcacag tgcccgaaaa aacgaaagtc aggaaacagc cgtgagcgat 1980
gtcagctgtg tgacgggatg ggacacaacg ctaaacagt taggaagcgg gatggcaacc 2040
agggccaacg cccaggaaga ggtctctctt cggggccgtg gcccgccct gagcagcctg 2100
cgtctcgtt agcgatgaca atggaacata aagatcgccc cttggttagg gtcattctga 2160
ctaactctg gagtcaccca gtcaaacaac gttcgggtga tatcaccgcg ctggttgact 2220
ccggagcgg catcactatt atttcggagg aggattggcc tactgattgg ccggtggtgg 2280
acaccgcgaa cccacagatc catggcatag gagggggaat tcccatgcga aaatcccggg 2340
atatgataga ggtgggggtt attaaccgag acgggtcgtt ggagcgaccc ctgctcctct 2400
tccccgcagt cgtatgggt agagggagta tcttaggaag agattgtctg cagggcctag 2460
ggtccgctt gacaaattta tagggagggc cactgttctc actgttgccg tacatctggc 2520
tattecgctc aaatggaagc cagaccgcac gcctgtgtgg attgaccagt ggccccctcc 2580
tgaaggtaaa cttgtaggcc taacgcaatt agtggaaaaa gaattacagt taggacatat 2640

agagccctca cttagtgtgtt ggaacacacc tgtttttcgt gatccggaag gcttccgggt 2700
cttatcgctt attgcatgat ttgcgcgtg ttaacgcaa gcttgccct tttggggccg 2760
tccaacaggg ggccaggtt ctctccgcgc tcccgctgg ctggccctg atggctctag 2820
acctcaagga ttgcttcttt tctatccctc ttgcggaaca agatcgcaa gcttttgcag 2880
ttacgctccc ctctgtgaat aaccaggccc ccgctcgaag attccaatgg aaggctcttg 2940
cccaagggat gacctgttct cccactatct gtcagttggg agtgggtcag gtgctcgagc 3000
ccttgcgact caagcaccca gctctgcgca tgttgcatat tatggacgat cttttgctag 3060
ccgcctcaag tcatgatggg ttggaagcgg caggggaagga gggtatcggg acattggaaa 3120
gagccgggtt cactatttct cgggataaga tccagagggg gcccgaggta caatatcttg 3180
ggtacaagtt aggcagtagc tatgtagcac ccgtaggctt ggtagcagaa cccaggatag 3240
ccacctgttg ggatgttcaa aagctgggtg ggtaacttca gtggcttcgc ccagcgtag 3300
ggatcccgcc acgactgatg ggtccctttt atgagcagtt acgaggggtc gatcctaacg 3360
aggcgagggg atggaatcta gacatgaaaa tggcctggag agagatcgta cagcttagca 3420
ctactgctgc cttggaacga tgggacctg cccagcctct ggaaggagcg gtcgctagat 3480
gtgaacaggg ggcaataggg gtccctgggac agggactgtc cacacacca aggccatgtt 3540
tgtggttatt ctccacccaa cccaccaagg cgtttactgc ttgggttagaa gtgctcacc 3600
ttttgattac taagctacgc gcttcggcag tgcgaacctt tggcaaggag gttgatatcc 3660
tcctgttgcc tgcagtcttc cgggaggacc tcccgctccc ggaggggagc ctgttagcac 3720
ttaggggggt tgcaggaaaa atcaggagta gtgacacgcc atctattttt gacattgcgc 3780
gtccactgca tgtttctctg aaagtgaggg ttaccgacca cctgtgccc ggaccactg 3840
tctttaccga cgcctcctca agcaccata aaggggtggg agtctggagg gagggcccaa 3900
gggtgggagat aaaagaaata gttgatttgg gggcaagtgt acaacaactg gaggcacgcg 3960
ctgtggccat ggcacttctg ctgtggccga caacgcccac taatgtagt actgactctg 4020
cgtttgttgc gaaaatgtta ctcaagatgg gacaggaggg agtcccgtct acagcggcg 4080
cttttatttt agaggatgcg ttaagccaaa ggtaagccat ggccgcccgt ctccacgtgc 4140
ggagtcattc tgaagtgccg gggtttttca cagaaggaaa tgacgtggca gatagccaag 4200
ccacctttca agcgtatccc ttgagagagg ctaaagatct tcataccgct ctccatattg 4260
gaccccgccg gctatccaaa gcggtgtaata tatctatgca gcaggctagg gaggttgttc 4320
agacctgccc gcattgtaat tcagccctg cgttggaggc cggggtaaac cctaggggtt 4380
tgggaccttc acagatatgg cagacagact ttacgcttga gcctagaatg gctccccgtt 4440
cctggctcgc tgttactgtg gacaccgct catcagcgat agtcgtaact cagcatggcc 4500
gtgttacatc gggtgctgca caacatcatt gggccacggc tatcgccgtt ttgggaagac 4560

caaaggccat aaaaacagat aacgggtcct gttcacgtc cagatccacg cgagagtggc 4620
tcgcgagatg ggggatagca cacaccacg ggattccggg aaattcccag ggtcaagcta 4680
tggtagagcg ggccaaccgg ctcttgaaag ataagatccg tgtgctcgcg gagggggacg 4740
gctttatgaa aagaatcccc accagcaaac aggggggaact attagccaag gcaatgtatg 4800
ccctcaatca ctttgagcgt ggtgaaaaca caaaaacacc gatacaaaaa cactggagac 4860
ctaccgttct tacagaagga cccccgggta aaatacgaat agagacaggg gagtgggaaa 4920
aaggatggaa cgtgctggtc tggggacgag gttatgccgc tgtgaaaaac agggacactg 4980
ataaggttat ttgggtaccc tctcggaag ttaaaccgga tgtcacccaa aaggatgagg 5040
tgactaagaa agatgaggcg agccctcttt ttgcaggcat ttctgactgg ataccctggg 5100
aagacgagca agaaggactc caaggagaaa ccgctagcaa caagcaagaa agaccgggag 5160
aagacaccct tgctgccaac gagagttaat tatattctca ttattggtgt cctggtcttg 5220
tgtgagggta cgggggtaag agctgatgtc cacttactcg agcagccagg gaacctttgg 5280
attacatggg ccaaccgtac aggccaaacg gatttttgcc tctctacaca gtcagccacc 5340
tccccctttc aaacatgttt gataggtatc ccgtccccta tttccgaggg tgattttaag 5400
ggatatgttt ctgatacaaa ttgcaccacc ttgggaactg atcggttagt ctcgtcagcc 5460
gactttactg gcggacctga caacagtacc accctcactt atcggaaggt ctcagtgttg 5520
ttgttaaagc tgaatgtctc tatgtgggat gagccacctg aactacagct gttaggttcc 5580
cagtctctcc ctaacattac taatattgct cagatttccg gtataaccgg gggatgcgta 5640
ggcttcagac cacaaggggt tccttggtat ctaggttggt ctagacagga ggccacgcgg 5700
tttctcctta gacaccctc tttctctaaa tccacggaac cgtttacagt ggtgacagcg 5760
gataggcaca atctttttat ggggagttag tactgcgggtg catatggcta cagattttgg 5820
aacatgtata actgctcaca ggtggggcgg cagtaccgt gtggtaatgc gcgcacgccc 5880
cgacagggtc ttctgaaat ccagtgtaca aggagaggag gcaaattgggt taatcaatca 5940
caggaaatta atgagtcgga gccgttcagc tttacggtga actgtacagc tagtagtttg 6000
ggtaatgcca gtgggtgttg cggaaaagca ggcacgattc tcccgggaaa gtgggtcgac 6060
agcacacaag gtagtttcac caaacaaaa gcgtaccac ccgcaatttt cctcatttgt 6120
ggggatcgcg catggcaagg aattcccagt cgtecggtag ggggcccctg ctatttaggc 6180
aagcttacca tgtagcacc taagcataca gatattctca aggtgcttgt caattcctcg 6240
cggacaggta taagacgtaa acgaagcacc tcacacctgg atgatacatg ctcagatgaa 6300
gtgcagcttt ggggtcctac agcaagaatc tttgcatcta tctagcccc ggggtagca 6360
gctgcgcaag ccttaagaga aattgagaga ctagcctgtt ggtccgttaa acaggctaac 6420
ttgacaacat cactcctcgg ggacttattg gatgatgtca cgagtattcg acacgcggtc 6480

ctgcagaacc gagcggtat tgacttcttg ctctagctc acggccatgg ctgtgaggac 6540
gttgccggaa tgtgctgttt caatttgagt gatcagagt agtctataca gaagaagttc 6600
cagctaataga aggaacatgt caataagatc ggcggtggata gcgacctaat tggaagttgg 6660
ctgcgaggac tattcggggg aataggagaa tgggcccgttc atttgctgaa aggactgctt 6720
ttggggccttg tagttatttt gttgctagta gtgtgcctgc cttgcctttt gcaaagtgtta 6780
tgcggtaata ggagaaagat gattaataac tccatcagct accacacgga atataagaag 6840
ctgcaaaagg cctgtgggca gcctgaaagc agaatagtat aaggcagtac atgggtggtg 6900
gtatagcgct tgcgagtcca tcgagcaagg caggaaagac agctattggt aattgtgaaa 6960
tacgcttttg tctgtgtgct gcaggagctg agctgactct gctggtggcc tcgctacca 7020
ctgtggcatc gatgcgatgt acgggccaga tatacgcgta tctgagggga ctagggtgtg 7080
tttaggcgaa aagcggggct tcggttgtag gcggttagga gtccccttag gatatagtag 7140
tttcgctttt gcatagggag ggggaaatgt agtcttatgc aatactcttg tagtcttgca 7200
acatggtaac gatgagttag caacatgcct tacaaggaga gaaaaagcac cgtgcatgcc 7260
gattggtgga agtaagggtg tacgatcgtg ccttattagg aaggcaacag acgggtctga 7320
catggattgg acgaaccact gaattccgca ttgcagagat attgtattta agtgcctagc 7380
tcgatacaat aaacgccatt tgaccattca ccacattggt gtgcacctgg gttgatggcc 7440
ggaccgttga ttccctgacg actacgagca cctgcatgaa gcagaaggct tcatttggtg 7500
accccgacgt gatagttagg gaatagtggt cggccacaga cggcgtggcg atcctgtctc 7560
catccgtctc gtctatcggg aggcgacttc gatgaccctg gtggaggggg ctgcggctta 7620
gggaggcaga agctgagtac cgtcggaggg gatccacagg acgggtgtgg tcgccatgat 7680
cgcgtagtcg atagtggctc caagtagcga agcgagcagg actgggcggc ggccaaagcg 7740
gtcggacagt gctccgagaa cgggtgcgca tagaaattgc atcaacgcat atagcgctag 7800
cagcacgcca tagtgactgg cgatgctgtc ggaatggacg atatcccgca agaggcccgg 7860
cagtaccggc ataaccaagc ctatgcctac agcatccagg gtgacggtgc cgaggatgac 7920
gatgagcgca ttgttagatt tcatacacgg tgctgactg cgttagcaat ttaactgtga 7980
taaactaccg cattaaagct ccaaacttgg ctgtttcctg tgtgaaattg ttatccgctc 8040
acaattccac acattatacg agccggaagc ataaagtgt aaacctgggg tgccaatga 8100
gtgagaattc ttgaagacga aagggcctcg tgatacgctt atttttatag gttaatgtca 8160
tgataataat ggtttcttag acgtcaggtg gcacttttcg gggaaatgtg cgcggaaccc 8220
ctatttgttt atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct 8280
gataaatgct tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg 8340
cccttattcc cttttttgcg gcattttgcc ttctgtttt tgctcaccca gaaacgctgg 8400

tgaaagtaaa agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc 8460
tcaacagcgg taagatcctt gagagttttc gccccgaaga acgttttcca atgatgagca 8520
cttttaaaagt tctgctatgt ggcgcggtat tatcccgtgt tgacgccggg caagagcaac 8580
tcggtcgccg catacactat tctcagaatg acttggttga gtactcacca gtcacagaaa 8640
agcatcttac ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg 8700
ataaactgct ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt 8760
ttttgcacaa catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg 8820
aagccatacc aaacgacgag cgtgacacca cgatgcctgc agcaatggca acaacgttgc 8880
gcaaactatt aactggcgaa ctacttactc tagcttcccc gcaacaatta atagactgga 8940
tgaggcgga taaagtgtca ggaccacttc tgcgctcggc ccttcgggct ggctggttta 9000
ttgctgataa atctggagcc ggtgagcgtg ggtctcgcgg tatcattgca gactggggc 9060
cagatggtaa gccctccgt atcgtagtta tctacacgac ggggagtcag gcaactatgg 9120
atgaacgaaa tagacagatc gctgagatag gtgcctcact gattaagcat tggtaactgt 9180
cagaccaagt ttactcatat atactttaga ttgatttaaa acttcatttt taatttataa 9240
ggatctaggt gaagatcctt tttgataatc tcatgaccaa aatcccttaa cgtgagtttt 9300
cgttccactg agcgtcagac cccgtagaaa agatcaaagg atcttcttga gatccttttt 9360
ttctgcgcgt aatctgctgc ttgcaaacaa aaaaaccacc gctaccagcg gtggtttgtt 9420
tgccggatca agagctacca actctttttc cgaaggtaac tggcttcagc agagcgcaga 9480
taccaaatac tgtccttcta gtgtagcgt agttaggcca ccacttcaag aactctgtag 9540
caccgcctac atacctcgt ctgctaatec tgttaccagt ggctgctgcc agtggcgata 9600
agtcgtgtct taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg 9660
gctgaacggg gggttcgtgc acacagccca gcttgagcg aacgacctac accgaactga 9720
gatacctaca gcgtgagcta tgagaaagcg ccacgcttc cgaagggaga aaggcggaca 9780
ggtatccggt aagcggcagg gtcggaacag gagagcgac gagggagctt ccagggggaa 9840
acgcctggta tctttatagt cctgtcgggt ttccgccact ctgacttgag cgtcgatttt 9900
tgtgatgctc gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac 9960
ggttccctggc cttttgctgg ccttttctc acatgttctt tcttgcgtta tccctgatt 10020
ctgtggataa ccgtattacc gcctttgagt gagctgatac cgctcgcgc agccgaacga 10080
ccgagcgcag cgagtcagtg agcgaggaag cggaagagcg cctgatgcgg tattttctcc 10140
ttacgcctct gtgcggtatt tcacacgcga tatggtgcac tctcagtaca atctgctctg 10200
atgccgcata gttaagccag tatacactcc gctatcgcta cgtgactggg tcatggctgc 10260
gccccgacac ccgccaacac ccgtgacgc gccctgacgg gcttgtctgc tcccgcatc 10320

WO 99/61590

PCT/US99/11780

9/18

cgcttacaga caagctgtga cegtcctccg gagctgcatg tgtcagaggt tttcacccgtc 10380
 atcacccgaaa cgcgcgaggg agctgcggta aagctcatca gcgtgggtcgt gaagcgattc 10440
 acagatgtct gcctgttcat cgcgctccag ctcgttgagt ttctccagaa gcgttaatgt 10500
 ctggcttctg ataaagcggg ccatgttaag ggcggttttt tctgttttg tcaattgatg 10560
 cctccgtgta agggggaatt tctgttcatg ggggtaataa taccgatgaa acgagagagg 10620
 atgctcacga tacgggttac tgatgatgaa catgcccggt tactggaacg ttgtgaggg 10680
 aaacaactgg cggatatgat gcggcgggac cagagaaaaa tcactcaggg tcaatgccag 10740
 cgcttcgtta atacagatgt aggtgttcca cagggtagcc agcagcatcc tgcgatgcag 10800
 atccggaaca taatggtgca gggcgctgac ttccgcgttt ccagacttta cgaaacacgg 10860
 aaaccgaaga ccattcatgt tgttgctcag gtcgcagacg ttttgcagca gcagtcgctt 10920
 cacgttcgct cgcgtatcgg tgattcatte tgctaaccag taaggcaacc ccgccagcct 10980
 agccgggtcc tcaacgacag gagcacgac atgagcacc gtggccagga cccaacgctg 11040
 cccgagatgc gccgcgtgcg gctgctggag atggcggacg cgatggatat gttctgcca 11100
 ggggttggtt gcgcattcac agttctccgc aagaattgat tggctccaat tcttgagtg 11160
 gtgaatccgt tagcgagggt ccgccggctt ccattcaggt cgagggtggc cggctccatg 11220
 caccgcgacg caacgcgggg aggcagacaa ggtataggg gcgatgcga tgtacgggcc 11280
 agatatacgc gtatctgagg ggactagggt gtgtttaggc gaaaagcggg gcttcgggtg 11340
 tacgcggtta ggagtcctct taggatatag tagtttcgct tttgcatagg gagggggaaa 11400
 tgtagtctta tgcaatactc ttgtagtctt gcaacatggt aacgatgagt tagcaacatg 11460
 ccttacaagg agagaaaaag caccgtgcat gccgattggt ggaagtaagg tggtagatc 11520
 gtgccttatt aggaaggcaa cagacgggtc tgacatggat tggacgaacc actgaattcc 11580
 gcattgcaga gatattgtat ttaagtcct agctcgatac aataaac 11627

<210> 2

<211> 1759

<212> DNA

<213> Chicken

<220>

<221> gene

<222> (1) .. (1759)

<223> chicken c-SRC cDNA

<220>

<221> CDS

<222> (112) .. (1710)

<400> 2

tctgacaccc atctgtctgt ctgtctgtgt gctgcaggag ctgagctgac tctgctgtgg 60

cctcgcgtac cactgtggcc aggcggtagc tgggacgtgc agcccaccac c atg ggg 117

WO 99/61590

PCT/US99/11780

10/18

Met Gly
1

| | |
|---|-----|
| agc agc aag agc aag ccc aag gac ccc agc cag cgc cgg cgc agc ctg | 165 |
| Ser Ser Lys Ser Lys Pro Lys Asp Pro Ser Gln Arg Arg Arg Ser Leu | |
| 5 10 15 | |
| gag cca ccc gac agc acc cac cac ggg gga ttc cca gcc tcg cag acc | 213 |
| Glu Pro Pro Asp Ser Thr His His Gly Gly Phe Pro Ala Ser Gln Thr | |
| 20 25 30 | |
| ccc aac aag aca gca gcc ccc gac acg cac cgc acc ccc agc cgc tcc | 261 |
| Pro Asn Lys Thr Ala Ala Pro Asp Thr His Arg Thr Pro Ser Arg Ser | |
| 35 40 45 50 | |
| ttt ggg acc gtg gcc acc gag ccc aag ctc ttc ggg ggc ttc aac act | 309 |
| Phe Gly Thr Val Ala Thr Glu Pro Lys Leu Phe Gly Gly Phe Asn Thr | |
| 55 60 65 | |
| tct gac acc gtt acg tcg ccg cag cgt gcc ggg gca ctg gct ggc ggc | 357 |
| Ser Asp Thr Val Thr Ser Pro Gln Arg Ala Gly Ala Leu Ala Gly Gly | |
| 70 75 80 | |
| gtc acc act ttc gtg gct ctc tac gac tac gag tcc cgg act gaa acg | 405 |
| Val Thr Thr Phe Val Ala Leu Tyr Asp Tyr Glu Ser Arg Thr Glu Thr | |
| 85 90 95 | |
| gac ttg tcc ttc aag aaa gga gaa cgc ctg cag att gtc aac aac acg | 453 |
| Asp Leu Ser Phe Lys Lys Gly Glu Arg Leu Gln Ile Val Asn Asn Thr | |
| 100 105 110 | |
| gaa ggt gac tgg tgg ctg gct cat tcc ctc act aca gga cag acg ggc | 501 |
| Glu Gly Asp Trp Trp Leu Ala His Ser Leu Thr Thr Gly Gln Thr Gly | |
| 115 120 125 130 | |
| tac atc ccc agt aac tat gtc gcg ccc tca gac tcc atc cag gct gaa | 549 |
| Tyr Ile Pro Ser Asn Tyr Val Ala Pro Ser Asp Ser Ile Gln Ala Glu | |
| 135 140 145 | |
| gag tgg tac ttt ggg aag atc act cgt cgg gag tcc gag cgg ctg ctg | 597 |
| Glu Trp Tyr Phe Gly Lys Ile Thr Arg Arg Glu Ser Glu Arg Leu Leu | |
| 150 155 160 | |
| ctc aac ccc gaa aac ccc cgg gga acc ttc ttg gtc cgg gag agc gag | 645 |
| Leu Asn Pro Glu Asn Pro Arg Gly Thr Phe Leu Val Arg Glu Ser Glu | |
| 165 170 175 | |
| acg aca aaa ggt gcc tat tgc ctc tcc gtt tct gac ttt gac aac gcc | 693 |
| Thr Thr Lys Gly Ala Tyr Cys Leu Ser Val Ser Asp Phe Asp Asn Ala | |
| 180 185 190 | |
| aag ggg ctc aat gtg aag cac tac aag atc cgc aag ctg gac agc ggc | 741 |
| Lys Gly Leu Asn Val Lys His Tyr Lys Ile Arg Lys Leu Asp Ser Gly | |
| 195 200 205 210 | |
| ggc ttc tac atc acc tca cgc aca cag ttc agc agc ctg cag cag ctg | 789 |
| Gly Phe Tyr Ile Thr Ser Arg Thr Gln Phe Ser Ser Leu Gln Gln Leu | |
| 215 220 225 | |
| gtg gcc tac tac tcc aaa cat gct gat ggc ttg tgc cac cgc ctg acc | 837 |
| Val Ala Tyr Tyr Ser Lys His Ala Asp Gly Leu Cys His Arg Leu Thr | |
| 230 235 240 | |
| aac gtc tgc ccc acg tcc aag ccc cag acc cag gga ctc gcc aag gac | 885 |

WO 99/61590

PCT/US99/11780

11/18

| Asn | Val | Cys | Pro | Thr | Ser | Lys | Pro | Gln | Thr | Gln | Gly | Leu | Ala | Lys | Asp | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|
| 245 | | | | | | | 250 | | | | | 255 | | | | |
| gcg | tgg | gaa | atc | ccc | cgg | gag | tcg | ctg | cgg | ctg | gag | gtg | aag | ctg | ggg | 933 |
| Ala | Trp | Glu | Ile | Pro | Arg | Glu | Ser | Leu | Arg | Leu | Glu | Val | Lys | Leu | Gly | |
| 260 | | | | | | 265 | | | | | 270 | | | | | |
| cag | ggc | tgc | ttt | gga | gag | gtc | tgg | atg | ggg | acc | tgg | aac | ggc | acc | acc | 981 |
| Gln | Gly | Cys | Phe | Gly | Glu | Val | Trp | Met | Gly | Thr | Trp | Asn | Gly | Thr | Thr | |
| 275 | | | | | 280 | | | | | 285 | | | | | 290 | |
| aga | gtg | gcc | ata | aag | act | ctg | aag | ccc | ggc | acc | atg | tcc | ccg | gag | gcc | 1029 |
| Arg | Val | Ala | Ile | Lys | Thr | Leu | Lys | Pro | Gly | Thr | Met | Ser | Pro | Glu | Ala | |
| | | | | 295 | | | | | 300 | | | | | 305 | | |
| ttc | ctg | cag | gaa | gcc | caa | gtg | atg | aag | aag | ctc | cgg | cat | gag | aag | ctg | 1077 |
| Phe | Leu | Gln | Glu | Ala | Gln | Val | Met | Lys | Lys | Leu | Arg | His | Glu | Lys | Leu | |
| | | | 310 | | | | | 315 | | | | | 320 | | | |
| gtt | cag | ctg | tac | gca | gtg | gtg | tcg | gaa | gag | ccc | atc | tac | atc | gtc | act | 1125 |
| Val | Gln | Leu | Tyr | Ala | Val | Val | Ser | Glu | Glu | Pro | Ile | Tyr | Ile | Val | Thr | |
| | | | 325 | | | | 330 | | | | | 335 | | | | |
| gag | tac | atg | agc | aag | ggg | agc | ctc | ctg | gat | ttc | ctg | aag | gga | gag | atg | 1173 |
| Glu | Tyr | Met | Ser | Lys | Gly | Ser | Leu | Leu | Asp | Phe | Leu | Lys | Gly | Glu | Met | |
| | 340 | | | | | 345 | | | | | 350 | | | | | |
| ggc | aag | tac | ctg | cgg | ctg | cca | cag | ctc | gtc | gat | atg | gct | gct | cag | att | 1221 |
| Gly | Lys | Tyr | Leu | Arg | Leu | Pro | Gln | Leu | Val | Asp | Met | Ala | Ala | Gln | Ile | |
| | 355 | | | | 360 | | | | | 365 | | | | | 370 | |
| gca | tcc | ggc | atg | gcc | tat | gtg | gag | agg | atg | aac | tac | gtg | cac | cga | gac | 1269 |
| Ala | Ser | Gly | Met | Ala | Tyr | Val | Glu | Arg | Met | Asn | Tyr | Val | His | Arg | Asp | |
| | | | | 375 | | | | | 380 | | | | | 385 | | |
| ctg | cgg | gcg | gcc | aac | atc | ctg | gtg | ggg | gag | aac | ctg | gtg | tgc | aag | gtg | 1317 |
| Leu | Arg | Ala | Ala | Asn | Ile | Leu | Val | Gly | Glu | Asn | Leu | Val | Cys | Lys | Val | |
| | | | 390 | | | | | 395 | | | | | 400 | | | |
| gct | gac | ttt | ggg | ctg | gca | cgc | ctc | atc | gag | gac | aac | gag | tac | aca | gca | 1365 |
| Ala | Asp | Phe | Gly | Leu | Ala | Arg | Leu | Ile | Glu | Asp | Asn | Glu | Tyr | Thr | Ala | |
| | | 405 | | | | | 410 | | | | | 415 | | | | |
| cgg | caa | ggt | gcc | aag | ttc | ccc | atc | aag | tgg | aca | gcc | ccc | gag | gca | gcc | 1413 |
| Arg | Gln | Gly | Ala | Lys | Phe | Pro | Ile | Lys | Trp | Thr | Ala | Pro | Glu | Ala | Ala | |
| | 420 | | | | | 425 | | | | | 430 | | | | | |
| ctc | tat | ggc | cgg | ttc | acc | atc | aag | tcg | gat | gtc | tgg | tcc | ttc | ggc | atc | 1461 |
| Leu | Tyr | Gly | Arg | Phe | Thr | Ile | Lys | Ser | Asp | Val | Trp | Ser | Phe | Gly | Ile | |
| | 435 | | | | 440 | | | | | 445 | | | | | 450 | |
| ctg | ctg | act | gag | ctg | acc | acc | aag | ggc | cgg | gtg | cca | tac | cca | ggg | atg | 1509 |
| Leu | Leu | Thr | Glu | Leu | Thr | Thr | Lys | Gly | Arg | Val | Pro | Tyr | Pro | Gly | Met | |
| | | | | 455 | | | | | 460 | | | | | 465 | | |
| gtc | aac | agg | gag | gtg | ctg | gac | cag | gtg | gag | agg | ggc | tac | cgc | atg | ccc | 1557 |
| Val | Asn | Arg | Glu | Val | Leu | Asp | Gln | Val | Glu | Arg | Gly | Tyr | Arg | Met | Pro | |
| | | | 470 | | | | 475 | | | | | | 480 | | | |
| tgc | ccg | ccc | gag | tgc | ccc | gag | tcg | ctg | cat | gac | ctc | atg | tgc | cag | tgc | 1605 |
| Cys | Pro | Pro | Glu | Cys | Pro | Glu | Ser | Leu | His | Asp | Leu | Met | Cys | Gln | Cys | |
| | | 485 | | | | | 490 | | | | | 495 | | | | |
| tgg | cgg | agg | gac | cct | gag | gag | cgg | ccc | act | ttt | gag | tac | ctg | cag | gcc | 1653 |

WO 99/61590

PCT/US99/11780

12/18

Trp Arg Arg Asp Pro Glu Glu Arg Pro Thr Phe Glu Tyr Leu Gln Ala
 500 505 510

ttc ctg gag gac tac ttc acc tcg aca gag ccc cag tac cag cct gga 1701
 Phe Leu Glu Asp Tyr Phe Thr Ser Thr Glu Pro Gln Tyr Gln Pro Gly
 515 520 525 530

gag aac cta taggcctgga gctcctcctg gaccagagga ctcgctgtgg ggtacaggg 1759
 Glu Asn Leu

<210> 3
 <211> 533
 <212> PRT
 <213> Chicken

<400> 3
 Met Gly Ser Ser Lys Ser Lys Pro Lys Asp Pro Ser Gln Arg Arg Arg
 1 5 10 15

Ser Leu Glu Pro Pro Asp Ser Thr His His Gly Gly Phe Pro Ala Ser
 20 25 30

Gln Thr Pro Asn Lys Thr Ala Ala Pro Asp Thr His Arg Thr Pro Ser
 35 40 45

Arg Ser Phe Gly Thr Val Ala Thr Glu Pro Lys Leu Phe Gly Gly Phe
 50 55 60

Asn Thr Ser Asp Thr Val Thr Ser Pro Gln Arg Ala Gly Ala Leu Ala
 65 70 75 80

Gly Gly Val Thr Thr Phe Val Ala Leu Tyr Asp Tyr Glu Ser Arg Thr
 85 90 95

Glu Thr Asp Leu Ser Phe Lys Lys Gly Glu Arg Leu Gln Ile Val Asn
 100 105 110

Asn Thr Glu Gly Asp Trp Trp Leu Ala His Ser Leu Thr Thr Gly Gln
 115 120 125

Thr Gly Tyr Ile Pro Ser Asn Tyr Val Ala Pro Ser Asp Ser Ile Gln
 130 135 140

Ala Glu Glu Trp Tyr Phe Gly Lys Ile Thr Arg Arg Glu Ser Glu Arg
 145 150 155 160

Leu Leu Leu Asn Pro Glu Asn Pro Arg Gly Thr Phe Leu Val Arg Glu
 165 170 175

Ser Glu Thr Thr Lys Gly Ala Tyr Cys Leu Ser Val Ser Asp Phe Asp
 180 185 190

Asn Ala Lys Gly Leu Asn Val Lys His Tyr Lys Ile Arg Lys Leu Asp
 195 200 205

Ser Gly Gly Phe Tyr Ile Thr Ser Arg Thr Gln Phe Ser Ser Leu Gln
 210 215 220

Gln Leu Val Ala Tyr Tyr Ser Lys His Ala Asp Gly Leu Cys His Arg
 225 230 235 240

Leu Thr Asn Val Cys Pro Thr Ser Lys Pro Gln Thr Gln Gly Leu Ala
 245 250 255

WO 99/61590

PCT/US99/11780

13/18

Lys Asp Ala Trp Glu Ile Pro Arg Glu Ser Leu Arg Leu Glu Val Lys
 260 265 270
 Leu Gly Gln Gly Cys Phe Gly Glu Val Trp Met Gly Thr Trp Asn Gly
 275 280 285
 Thr Thr Arg Val Ala Ile Lys Thr Leu Lys Pro Gly Thr Met Ser Pro
 290 295 300
 Glu Ala Phe Leu Gln Glu Ala Gln Val Met Lys Lys Leu Arg His Glu
 305 310 315 320
 Lys Leu Val Gln Leu Tyr Ala Val Val Ser Glu Glu Pro Ile Tyr Ile
 325 330 335
 Val Thr Glu Tyr Met Ser Lys Gly Ser Leu Leu Asp Phe Leu Lys Gly
 340 345 350
 Glu Met Gly Lys Tyr Leu Arg Leu Pro Gln Leu Val Asp Met Ala Ala
 355 360 365
 Gln Ile Ala Ser Gly Met Ala Tyr Val Glu Arg Met Asn Tyr Val His
 370 375 380
 Arg Asp Leu Arg Ala Ala Asn Ile Leu Val Gly Glu Asn Leu Val Cys
 385 390 395 400
 Lys Val Ala Asp Phe Gly Leu Ala Arg Leu Ile Glu Asp Asn Glu Tyr
 405 410 415
 Thr Ala Arg Gln Gly Ala Lys Phe Pro Ile Lys Trp Thr Ala Pro Glu
 420 425 430
 Ala Ala Leu Tyr Gly Arg Phe Thr Ile Lys Ser Asp Val Trp Ser Phe
 435 440 445
 Gly Ile Leu Leu Thr Glu Leu Thr Thr Lys Gly Arg Val Pro Tyr Pro
 450 455 460
 Gly Met Val Asn Arg Glu Val Leu Asp Gln Val Glu Arg Gly Tyr Arg
 465 470 475 480
 Met Pro Cys Pro Pro Glu Cys Pro Glu Ser Leu His Asp Leu Met Cys
 485 490 495
 Gln Cys Trp Arg Arg Asp Pro Glu Glu Arg Pro Thr Phe Glu Tyr Leu
 500 505 510
 Gln Ala Phe Leu Glu Asp Tyr Phe Thr Ser Thr Glu Pro Gln Tyr Gln
 515 520 525
 Pro Gly Glu Asn Leu
 530

<210> 4
 <211> 2187
 <212> DNA
 <213> Homo sapiens

<220>
 <221> gene
 <222> (1)..(2187)

WO 99/61590

PCT/US99/11780

14/18

<223> human c-SRC cDNA

<220>

<221> CDS

<222> (134) .. (1483)

<400> 4

```

gcgccgcgctc ccgcaggccg tgatgccgcc cgcgcggagg tggcccggac cgcagtgcc 60
caagagagct ctaatggtac caagtgcacag gttggettta ctgtgactcg gggacgccag 120
agctcctgag aag atg tca gca ata cag gcc gcc tgg cca tcc ggt aca 169
      Met Ser Ala Ile Gln Ala Ala Trp Pro Ser Gly Thr
        1             5             10

gaa tgt att gcc aag tac aac ttc cac ggc act gcc gag cag gac ctg 217
Glu Cys Ile Ala Lys Tyr Asn Phe His Gly Thr Ala Glu Gln Asp Leu
      15             20             25

ccc ttc tgc aaa gga gac gtg ctc acc att gtg gcc gtc acc aag gac 265
Pro Phe Cys Lys Gly Asp Val Leu Thr Ile Val Ala Val Thr Lys Asp
      30             35             40

ccc aac tgg tac aaa gcc aaa aac aag gtg ggc cgt gag ggc atc atc 313
Pro Asn Trp Tyr Lys Ala Lys Asn Lys Val Gly Arg Glu Gly Ile Ile
      45             50             55             60

cca gcc aac tac gtc cag aag cgg gag ggc gtg aag gcg ggt acc aaa 361
Pro Ala Asn Tyr Val Gln Lys Arg Glu Gly Val Lys Ala Gly Thr Lys
      65             70             75

ctc agc ctc atg cct tgg ttc cac ggc aag atc aca cgg gag cag gct 409
Leu Ser Leu Met Pro Trp Phe His Gly Lys Ile Thr Arg Glu Gln Ala
      80             85             90

gag cgg ctt ctg tac ccg ccg gag aca ggc ctg ttc ctg gtg cgg gag 457
Glu Arg Leu Leu Tyr Pro Pro Glu Thr Gly Leu Phe Leu Val Arg Glu
      95             100             105

agc acc aac tac ccc gga gac tac acg ctg tgc gtg agc tgc gac ggc 505
Ser Thr Asn Tyr Pro Gly Asp Tyr Thr Leu Cys Val Ser Cys Asp Gly
      110             115             120

aag gtg gag cac tac cgc atc atg tac cat gcc agc aag ctc agc atc 553
Lys Val Glu His Tyr Arg Ile Met Tyr His Ala Ser Lys Leu Ser Ile
      125             130             135             140

gac gag gag gtg tac ttt gag aac ctc atg cag ctg gtg gag cac tac 601
Asp Glu Glu Val Tyr Phe Glu Asn Leu Met Gln Leu Val Glu His Tyr
      145             150             155

acc tca gac gca gat gga ctc tgt acg cgc ctc att aaa cca aag gtc 649
Thr Ser Asp Ala Asp Gly Leu Cys Thr Arg Leu Ile Lys Pro Lys Val
      160             165             170

atg gag ggc aca gtg gcg gcc cag gat gag ttc tac cgc agc ggc tgg 697
Met Glu Gly Thr Val Ala Ala Gln Asp Glu Phe Tyr Arg Ser Gly Trp
      175             180             185

gcc ctg aac atg aag gag ctg aag ctg ctg cag acc atc ggg aag ggg 745
Ala Leu Asn Met Lys Glu Leu Lys Leu Leu Gln Thr Ile Gly Lys Gly
      190             195             200

gag ttc gga gac gtg atg ctg ggc gat tac cga ggg aac aaa gtc gcc 793

```

[illegible]

WO 99/61590

PCT/US99/11780

16/18

tgaactgagc cccagcgggc tggcgggcct ttttcctgcg tcccagcctg caccctcccg 1633
 gccccgtctc tcttggaacc acctgtgggg cctggggagc ccactgaggg gccagggagg 1693
 aaggaggcca cggagcggga ggcagcgcgc caccacgtcg ggcttccctg gcctcccgcc 1753
 actcgccttc ttagagtttt attcctttcc ttttttgaga ttttttttcc gtgtgtttat 1813
 tttttattat ttttcaagat aaggagaaaag aaagtaccca gcaaattgggc attttacaag 1873
 aagtacgaat cttatttttc ctgtcctgcc cgtgaggggtg gggggggaccg ggccccctctc 1933
 tagggacccc tcgccccagc ctcattcccc attctgtgtc ccatgtcccg tgtctcctcg 1993
 gtcgccccgt gtttgcgctt gaccatgttg cactgtttgc atgcgcccga ggcagacgtc 2053
 tgtcaggggc ttggatttcg tgtgcgcgtg ccaccgcgcc acccgccctg tgagatggaa 2113
 ttgtaataaa ccacgccatg aggacaccgc cgccgcctc ggcgcttcc caccgaaaa 2173
 aaaaaaaaaa aaaa 2187

<210> 5
 <211> 450
 <212> PRT
 <213> Homo sapiens

<400> 5
 Met Ser Ala Ile Gln Ala Ala Trp Pro Ser Gly Thr Glu Cys Ile Ala
 1 5 10 15
 Lys Tyr Asn Phe His Gly Thr Ala Glu Gln Asp Leu Pro Phe Cys Lys
 20 25 30
 Gly Asp Val Leu Thr Ile Val Ala Val Thr Lys Asp Pro Asn Trp Tyr
 35 40 45
 Lys Ala Lys Asn Lys Val Gly Arg Glu Gly Ile Ile Pro Ala Asn Tyr
 50 55 60
 Val Gln Lys Arg Glu Gly Val Lys Ala Gly Thr Lys Leu Ser Leu Met
 65 70 75 80
 Pro Trp Phe His Gly Lys Ile Thr Arg Glu Gln Ala Glu Arg Leu Leu
 85 90 95
 Tyr Pro Pro Glu Thr Gly Leu Phe Leu Val Arg Glu Ser Thr Asn Tyr
 100 105 110
 Pro Gly Asp Tyr Thr Leu Cys Val Ser Cys Asp Gly Lys Val Glu His
 115 120 125
 Tyr Arg Ile Met Tyr His Ala Ser Lys Leu Ser Ile Asp Glu Glu Val
 130 135 140
 Tyr Phe Glu Asn Leu Met Gln Leu Val Glu His Tyr Thr Ser Asp Ala
 145 150 155 160
 Asp Gly Leu Cys Thr Arg Leu Ile Lys Pro Lys Val Met Glu Gly Thr
 165 170 175
 Val Ala Ala Gln Asp Glu Phe Tyr Arg Ser Gly Trp Ala Leu Asn Met

WO 99/61590

PCT/US99/11780

17/18

| 180 | | | | | | | | | | 185 | | | | | 190 | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|--|--|
| Lys | Glu | Leu | Lys | Leu | Leu | Gln | Thr | Ile | Gly | Lys | Gly | Glu | Phe | Gly | Asp | | | | |
| | | 195 | | | | | 200 | | | | | 205 | | | | | | | |
| Val | Met | Leu | Gly | Asp | Tyr | Arg | Gly | Asn | Lys | Val | Ala | Val | Lys | Cys | Ile | | | | |
| | 210 | | | | | 215 | | | | | 220 | | | | | | | | |
| Lys | Asn | Asp | Ala | Thr | Ala | Gln | Ala | Phe | Leu | Ala | Glu | Ala | Ser | Val | Met | | | | |
| | 225 | | | | | 230 | | | | | 235 | | | | 240 | | | | |
| Thr | Gln | Leu | Arg | His | Ser | Asn | Leu | Val | Gln | Leu | Leu | Gly | Val | Ile | Val | | | | |
| | | | | 245 | | | | | 250 | | | | | 255 | | | | | |
| Glu | Glu | Lys | Gly | Gly | Leu | Tyr | Ile | Val | Thr | Glu | Tyr | Met | Ala | Lys | Gly | | | | |
| | | | 260 | | | | | 265 | | | | | 270 | | | | | | |
| Ser | Leu | Val | Asp | Tyr | Leu | Arg | Ser | Arg | Gly | Arg | Ser | Val | Leu | Gly | Gly | | | | |
| | | 275 | | | | | 280 | | | | | 285 | | | | | | | |
| Asp | Cys | Leu | Leu | Lys | Phe | Ser | Leu | Asp | Val | Cys | Glu | Ala | Met | Glu | Tyr | | | | |
| | 290 | | | | | 295 | | | | | 300 | | | | | | | | |
| Leu | Glu | Gly | Asn | Asn | Phe | Val | His | Arg | Asp | Leu | Ala | Ala | Arg | Asn | Val | | | | |
| | 305 | | | | 310 | | | | | | 315 | | | | 320 | | | | |
| Leu | Val | Ser | Glu | Asp | Asn | Val | Ala | Lys | Val | Ser | Asp | Phe | Gly | Leu | Thr | | | | |
| | | | | 325 | | | | | 330 | | | | | 335 | | | | | |
| Lys | Glu | Ala | Ser | Ser | Thr | Gln | Asp | Thr | Gly | Lys | Leu | Pro | Val | Lys | Trp | | | | |
| | | | 340 | | | | | 345 | | | | | 350 | | | | | | |
| Thr | Ala | Pro | Glu | Ala | Leu | Arg | Glu | Lys | Lys | Phe | Ser | Thr | Lys | Ser | Asp | | | | |
| | | 355 | | | | | 360 | | | | | 365 | | | | | | | |
| Val | Trp | Ser | Phe | Gly | Ile | Leu | Leu | Trp | Glu | Ile | Tyr | Ser | Phe | Gly | Arg | | | | |
| | 370 | | | | | 375 | | | | | 380 | | | | | | | | |
| Val | Pro | Tyr | Pro | Arg | Ile | Pro | Leu | Lys | Asp | Val | Val | Pro | Arg | Val | Glu | | | | |
| | 385 | | | | 390 | | | | | | 395 | | | | 400 | | | | |
| Lys | Gly | Tyr | Lys | Met | Asp | Ala | Pro | Asp | Gly | Cys | Pro | Pro | Ala | Val | Tyr | | | | |
| | | | | 405 | | | | | 410 | | | | | 415 | | | | | |
| Glu | Val | Met | Lys | Asn | Cys | Trp | His | Leu | Asp | Ala | Ala | Met | Arg | Pro | Ser | | | | |
| | | | 420 | | | | | 425 | | | | | 430 | | | | | | |
| Phe | Leu | Gln | Leu | Arg | Glu | Gln | Leu | Glu | His | Ile | Lys | Thr | His | Glu | Leu | | | | |
| | | 435 | | | | | 440 | | | | | 445 | | | | | | | |
| His | Leu | | | | | | | | | | | | | | | | | | |
| | 450 | | | | | | | | | | | | | | | | | | |

<210> 6

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:9E10-myc
epitope tag

<400> 6

WO 99/61590

PCT/US99/11780

18/18

Val Asp Met Glu Gln Lys Leu Ile Ala Glu Glu Asp Leu Asn
1 5 10